

Name: Brendon Charles

Teacher/Class: Ms. Berios

SYDNEY TECHNICAL HIGH SCHOOL

YEAR 12

HSC ASSESSMENT TASK 2

MARCH 2006

EXTENSION 1 MATHEMATICS

Time Allowed: 70 minutes

Instructions:

- Write your name and class at the top of each page.
- All necessary working must be shown. Marks may be deducted for careless or badly arranged work.
- Marks indicated are a guide only and may be varied if necessary.
- Start each question on a new page.
- Standard integrals can be found on the last page.


Question 1	Question 2	Question 3	Question 4	Question 5	Question 6	Total
/10	/10	/10	/10	/10	/10	/60

Question 1 (10 marks)

Marks

- a) Find the exact value of
- i. $\tan\left(\frac{2\pi}{3}\right)$ 1
 - ii. $\sin\left(-\frac{\pi}{3}\right)$ 1
- b) Find
- $$\lim_{x \rightarrow 0} \frac{\sin 2x}{x}$$
- 1
- c) Given that $\int_1^5 f(x) dx = 4$ find the value of k 3
 for which $\int_1^5 [f(x) + kx] dx = 28$.
- d) A river 60m wide is surveyed for its depth every 10m across its width.

The depth at each point surveyed is shown on the diagram.




- i. Find the cross-sectional area of the river using Simpsons Rule 3
- ii. Hence find the volume of water passing this point per second if the water flows at 5m/s. 1

Question 2 (10 marks) Start a new page

Marks

- a) The point of intersection of $y = x^3$ and $y = 2 - x$ is the point A (1, 1)

3



Find the shaded area.

- b) If $y = \sin 2x^\circ$


i. Express $2x^\circ$ in radian measure

1

ii. Find $\int \sin 2x^\circ dx$

2

- c)



i. Find x_1

1

ii. Calculate the volume generated when the shaded region (shown above) between the line $y = x$, the circle $x^2 + y^2 = 2$ and the x axis is rotated around the x axis.

3

Question 3 (10 marks) Start a new page


a) If $y = a \cos nx + b \sin nx$

show that $\frac{d^2y}{dx^2} + n^2y = 0$ 3

b) i. Differentiate $x\sqrt{x+3}$ and simplify your answer as far as possible. 2

ii. Hence find $\int \frac{x+2}{\sqrt{x+3}} dx$ 1

c) The sector below has area of 25cm^2 . It is contained in a circle of radius r cm and the arc AB subtends an angle at the centre of the circle of θ radians.



i. Show the perimeter of the sector is given by $P = 2r + \frac{50}{r}$ 1

ii. Find r for which the perimeter is a minimum. 3

Question 4 (10 marks) Start a new page

- a) i. Sketch $y = 3 \cos 2x$ for $0 \leq x \leq \pi$ 2
- ii. Find the area enclosed by $y = 3 \cos 2x$, the x axis, $x = 0$
and $x = \frac{\pi}{2}$ 3
- b) i. Express $\sin x + \sqrt{3} \cos x$ in the form $A \sin(x + \theta)$ for $0 < \theta < \frac{\pi}{2}$ 2
- ii. Hence solve $\sin x + \sqrt{3} \cos x = \sqrt{2}$ for $0 \leq x \leq 2\pi$. 3

Question 5 (10 marks) Start a new page

- a) Find $\int \frac{\cos \sqrt{x}}{\sqrt{x}} dx$ using the substitution $u = \sqrt{x}$ 3
- b) Find $\int \frac{x}{\sqrt{1-x}} dx$ using the substitution $u = 1-x$. 4
- c) Find $\int \cos^2 3x dx$ 3

Question 6 (10 marks) Start a new page

- a) Evaluate $\int_0^1 \frac{x}{(x^2 + 2)^2} dx$ using the substitution $u = x^2 + 2$ 3
- b) i. Prove $\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$ 1
- ii. Hence or otherwise evaluate $\int_0^{\pi/6} \sin 4x \cdot \cos 2x dx$ 3
- c) i. Sketch $y = 2^x$ 1
- ii. If n is a positive integer, by considering the graph of $y = 2^x$ 2
- show that $2^n < \int_n^{n+1} 2^x dx < 2 \cdot 2^n$

Question 1

a) i) $\tan\left(\frac{2\pi}{3}\right) = \tan\left(\pi - \frac{\pi}{3}\right)$
 $= -\tan\frac{\pi}{3}$
 $= -\sqrt{3}$

ii) $\sin\left(-\frac{\pi}{3}\right) = \sin\left(2\pi - \frac{\pi}{3}\right)$
 $= -\sin\frac{\pi}{3}$
 $= -\frac{\sqrt{3}}{2}$

b) $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = \lim_{x \rightarrow 0} \frac{\sin 2x \cdot 2}{2x}$
 $= 2$

c) $\int_1^5 [f(x) + kx] dx = \int_1^5 f(x) dx + \int_1^5 kx dx$
 $4 + \left[\frac{kx^2}{2}\right]_1^5 = 28$
 $\frac{25k}{2} - \frac{k}{2} = 24$
 $24k = 48$
 $k = 2$

d) i) $A = \frac{10}{3}(0+0+4(5+15+7)+2(12+14))$
 $A = 533\frac{1}{3} m^2$

ii) $V = 533\frac{1}{3} \times 5$
 $= 2666\frac{2}{3} m^3$

Question 2

a) $A = \int_0^4 x^3 dx + \frac{(1 \times 1)}{2} + \frac{(2 \times 2)}{2}$
 $= \left[\frac{x^4}{4}\right]_0^4 + 2\frac{1}{2}$
 $A = 2\frac{3}{4} \text{ unit}^2$

b) $\pi^c = 180^\circ$

i) $\therefore 2x^\circ = \frac{2 \times \pi \times c}{180}$
 $= \frac{\pi x}{90} \text{ radians}$

ii) $\int \sin 2x^\circ dx = \int \sin \frac{\pi}{90} x dx$
 $= -\frac{90}{\pi} \cos \frac{\pi x}{90} + c$

c) i) sineq. $y=x$ $x^2 + y^2 = 2$
 $x^2 + x^2 = 2$
 $2x^2 = 2$
 $x_1 = 1$

ii) $V = \pi \int_0^1 x^2 dx + \pi \int_1^{\sqrt{2}} (2-x^2) dx$
 $= \pi \left\{ \left[\frac{x^3}{3}\right]_0^1 + \left[2x - \frac{x^3}{3}\right]_1^{\sqrt{2}} \right\}$
 $= \pi \left[\frac{1}{3} + \left(2\sqrt{2} - \frac{2\sqrt{2}}{3}\right) - \left(2 - \frac{1}{3}\right) \right]$
 $= \pi \left[-\frac{4}{3} + \frac{6\sqrt{2} - 2\sqrt{2}}{3} \right]$
 $= \pi \left[\frac{4\sqrt{2} - 4}{3} \right]$

Question 3

a) $y = a \cos nx + b \sin nx$
 $\frac{dy}{dx} = -a n \sin nx + b n \cos nx$
 $\frac{d^2y}{dx^2} = -a n^2 \cos nx - b n^2 \sin nx$
 sub into $\frac{d^2y}{dx^2} + n^2 y = 0$
 LHS = $-a n^2 \cos nx - b n^2 \sin nx + n^2(a \cos nx + b \sin nx)$
 $= 0$
 $= \text{RHS}$

b) i) $y = a \sqrt{x+3}$
 Let $u = x$ $v = \sqrt{x+3} = (x+3)^{1/2}$
 $u' = 1$ $v' = \frac{1}{2}(x+3)^{-1/2} = \frac{1}{2\sqrt{x+3}}$
 $\therefore \frac{dy}{dx} = \sqrt{x+3} + \frac{xc}{2\sqrt{x+3}}$
 $= \frac{2(x+3) + xc}{2\sqrt{x+3}}$
 $= \frac{3xc + 6}{2\sqrt{x+3}}$
 $\frac{dy}{dx} = \frac{3[xc+2]}{2\sqrt{x+3}}$

ii) $\therefore \int \frac{x+2}{\sqrt{x+3}} dx = \frac{2}{3} x \sqrt{x+3} + c$


c) i) $P = 2r + \text{arc length AB}$
 $= 2r + r\theta$ — (1)
 since $\frac{1}{2} r^2 \theta = 25$
 $\theta = \frac{50}{r^2}$ sub into (1)
 $\therefore P = 2r + r \left[\frac{50}{r^2} \right]$
 $P = 2r + \frac{50}{r} = 2r + 50r^{-1}$

ii) $\frac{dP}{dr} = 2 - 50r^{-2}$
 $\frac{d^2P}{dr^2} = 100r^{-3}$
 stpts $2 - \frac{50}{r^2} = 0$
 $2r^2 = 50$
 $r = \pm 5$ $r > 0 \therefore r = 5$

test max/min
 if $r = 5$ $\frac{d^2P}{dr^2} > 0 \therefore \text{min}$
 $\therefore \text{min Perimeter if } r = 5 \text{ cm}$

Question 4

a) i) amplitude = 3 period $\frac{2\pi}{2} = \pi$



ii) $A = 2 \int_0^{\pi/4} 3 \cos 2x dx$
 $= 6 \left[\frac{1}{2} \sin 2x \right]_0^{\pi/4}$
 $= 3 \left[\sin \frac{\pi}{2} - \sin 0 \right]$
 $= 3 \text{ unit}^2$

b) i) $A = \sqrt{1+3} \therefore A = 2$
 $2 \left[\frac{1}{2} \sin x + \frac{\sqrt{3}}{2} \cos x \right]$
 $= A \sin(x + \theta)$
 $\cos \theta = \frac{1}{2}, \sin \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = \frac{\pi}{3}$
 $\therefore \sin x + \sqrt{3} \cos x = 2 \sin\left(x + \frac{\pi}{3}\right)$

ii) $2 \sin\left(x + \frac{\pi}{3}\right) = \sqrt{2}$ $\sqrt{2} \mid \text{A} \sqrt{2}$
 $\sin\left(x + \frac{\pi}{3}\right) = \frac{\sqrt{2}}{2}$ $\frac{\sqrt{2}}{2} \mid \text{A} \sqrt{2}$
 $x + \frac{\pi}{3} = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$
 $\therefore x = \frac{5\pi}{12}, \frac{23\pi}{12}$

Question 5

a) $u = \sqrt{x} = x^{1/2}$
 $\frac{du}{dx} = \frac{1}{2} x^{-1/2} = \frac{1}{2\sqrt{x}}$

$du = \frac{dx}{2\sqrt{x}}$

$\therefore dx = 2\sqrt{x} du$

$\int \frac{\cos \sqrt{x}}{\sqrt{x}} dx = \int \frac{\cos u \cdot 2\sqrt{x} du}{\sqrt{x}}$

$= 2 \int \cos u du$

$= 2 \sin u + C$

$= 2 \sin \sqrt{x} + C$

b) $u = 1-x$

$\frac{du}{dx} = -1$

$-du = dx$

$\int \frac{x}{\sqrt{1-x}} dx = \int \frac{1-u}{\sqrt{u}} \cdot -du$

$= - \int (1-u) u^{-1/2} du$

$= - \int (u^{-1/2} - u^{1/2}) du$

$= - \left[\frac{u^{1/2}}{1/2} - \frac{u^{3/2}}{3/2} \right]$

$= - 2\sqrt{1-x} + \frac{2}{3}(1-x)^{3/2} + C$

c) $\cos 2\theta = 2\cos^2 \theta - 1$

$\therefore \int \cos^2 3x dx = \frac{1}{2} \int (\cos 6x + 1) dx$

$= \frac{1}{2} \left[\frac{1}{6} \sin 6x + x \right] + C$

$= \frac{1}{12} \sin 6x + \frac{x}{2} + C$

Question 6

a) $u = x^2 + 2$ $x=1 \rightarrow u=3$

$\frac{du}{dx} = 2x$ $x=0 \rightarrow u=2$

$dx = \frac{du}{2x}$

$\int_0^1 \frac{x}{(x^2+2)^2} dx = \int_2^3 \frac{\frac{du}{2x}}{u^2} \cdot \frac{du}{2x}$

$= \frac{1}{2} \int_2^3 u^{-2} du$

$= \frac{1}{2} \left[-\frac{1}{u} \right]_2^3$

$= \frac{1}{12}$

b) i) LHS = $\sin(A+B) + \sin(A-B)$

$= \sin A \cos B + \cos A \sin B + \sin A \cos B - \cos A \sin B$

$= 2 \sin A \cdot \cos B$

$= \text{RHS}$

ii) $\sin 4x \cdot \cos 2x = \frac{1}{2} [\sin 6x + \sin 2x]$


$\frac{1}{2} \int_0^{\pi/6} (\sin 6x + \sin 2x) dx$

$= \frac{1}{2} \left[-\frac{1}{6} \cos 6x - \frac{1}{2} \cos 2x \right]_0^{\pi/6}$

$= \frac{1}{2} \left[-\frac{1}{6} \cos \pi - \frac{1}{2} \cos \frac{\pi}{3} - \left(-\frac{1}{6} - \frac{1}{2} \right) \right]$

$= \frac{1}{2} \left[\frac{1}{6} - \frac{1}{4} + \frac{1}{6} + \frac{1}{2} \right] = \frac{7}{24}$

c) i)



ii) area lower rectangle $< \int_n^{n+1} 2^x dx < \text{area upper rectangle}$

$2^n \times 1 < \int_n^{n+1} 2^x dx < 2^{n+1} \times 1$

$2^n < \int_n^{n+1} 2^x dx < 2 \cdot 2^n$