

SYDNEY TECHNICAL HIGH SCHOOL

YEAR 12 ASSESSMENT TASK 3

JUNE 2003

MATHEMATICS

Time Allowed: 70 minutes

Instructions:

- * Attempt all questions.
- * Answers to be written on the paper provided.
- * Start each question on a new page.
- * All necessary working should be shown.
- * Marks may not be awarded for careless or badly arranged working.
- * This question paper must be stapled on top of your answers.
- * Marks shown are for guidance and may be changed slightly if needed.
- * Standard integrals are attached and may be removed for your convenience.

Name: Hassan Salem

Teacher: Hassan Salem

Question 1	Question 2	Question 3	Question 4	Question 5	Total
12 /12	11 /12	9 /13	10 /11	12 /12	60


54

Question 1 (12 marks)

Marks

- a) Change 2.3 radians to degrees and minutes 1
- b) Find $\tan 3.2$ to 3 decimal places 1
- c) Find the exact value of
- i) $\sin \frac{3\pi}{4}$ 1
- ii) $\cos \left(\frac{-\pi}{6} \right)$ 1
- d) Solve $2\sin x = -\sqrt{3}$ for $0 \leq x \leq 2\pi$ 2
- e) AC and BD are the arcs of two concentric circles with centre O.
 $AB = OB = CD = OD = 10\text{cm}$.

$$\angle AOC = \frac{\pi}{15}$$



- Find i) the perimeter of Sector OAC 2
in terms of π
- ii) the area of ABDC in terms 2
of π

- f) If $f(x) = 2\cos 2x$ find $f' \left(\frac{\pi}{4} \right)$ 2

Question 2 (12 marks)

- a) Find
- i) $\frac{d}{dx}(\sin x^2)$ 1
- ii) $\frac{d}{dx}(\tan^2 x)$ 2
- b) The gradient function of a curve is given by $\frac{dy}{dx} = 6\cos 2x - \sin x$.
If the curve passes through the origin, find the equation of the curve 3
- c) If $y = 3\sin x - 4\cos x$ show that $\frac{d^2y}{dx^2} + y = 0$ 3

d) Find $\int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \cos 2x dx$ 3

Question 3 (13 marks)

a) Evaluate $\int_0^2 (1-2x)^3 dx$ 3

b) A curve is defined as follows

$$f(x) = \begin{cases} -1 & \text{for } -2 \leq x < 0 \\ x^2 - 1 & \text{for } 0 \leq x \leq 3 \end{cases}$$

i) Sketch the curve neatly showing a suitable scale on both axes 2

ii) State the co-ordinates of the point where the curve cuts the x axis. 1

iii) Find the area between the curve, the x axis and the lines $x = -2$ and $x = 3$. 4

c) Solve $2 \sin^2 x - \sin x = 0$ for $0 \leq x \leq 2\pi$ 3

Question 4 (11 marks)

a) i) Sketch $y = 4 \sin \frac{x}{2}$ in the domain $-2\pi \leq x \leq 2\pi$. State its period and amplitude 3

ii) Hence find the number of solutions to the equation $\sin \frac{x}{2} = \frac{1}{4}$ in the domain $-2\pi \leq x \leq 2\pi$ 1

b) i) The table below has been completed for $f(x) = \sqrt{\frac{x^2-1}{x}}$

x	1	1.25	1.5	1.75	2
f(x)	0	0.67	0.91	1.09	1.22

1st 4 2 4 Last

ii) By using Simpson's Rule and the five function values in the table find $\int_1^2 f(x) dx$ 2

- c) The area bounded by the curve $y = \sqrt{x^3}$, the x axis and the line $x = 3$, is rotated around the x axis. Find the volume of the solid formed (leave your answer in terms of π).

3

- d) Differentiate $y = x \sin 2x$

2

Question 5 (12 marks)


- a) i) Find the x values of the points of intersection of the line $y = 2x$ and the curve $3y = x^2$ 1
 ii) On the same axes sketch $y = 2x$ and $3y = x^2$, showing clearly the points of intersection. Label the line and the curve and shade the enclosed area. 2
 iii) Find the shaded area 2

- b) A metal gutter open at the top and ends, is bent up from material 30cm wide to form a rectangular cross section.

- i) If the base of the gutter is x cm wide, find the height of the gutter. 1
 ii) Show that the area of the gutter cross section is given by

$$A = 15x - \frac{x^2}{2} \quad 1$$

- iii) Find the value of x for which A is a maximum. 3



- c) Find $\int \tan^2 \theta d\theta$

2

Question 1 (12 mks)

1) $2.3 \text{ radians} = \frac{131^\circ 47'}{1}$ (1)

2) $\tan 3.2 = \frac{0.058}{1}$ (1)


3) i) $\sin \frac{3\pi}{4} = \sin(\pi - \frac{\pi}{4})$
 $= \sin \frac{\pi}{4}$
 $= \frac{1}{\sqrt{2}}$ (1)

ii) $\cos(-\frac{\pi}{6}) = \cos \frac{\pi}{6}$
 $= \frac{\sqrt{3}}{2}$ (1)

4) $2 \sin x = -\sqrt{3}$
 $\sin x = -\frac{\sqrt{3}}{2}$

S	A
√T	CV

 acute $x = \frac{\pi}{3}$
 $\therefore x = \frac{4\pi}{3}, \frac{5\pi}{3}$ (2)



i) $P = 40 + 20 \times \frac{\pi}{15}$
 $P = (40 + \frac{4\pi}{3}) \text{ cm}$ (2)

ii) $A = (\frac{1}{2} \times 20^2 \times \frac{\pi}{15}) -$
 $(\frac{1}{2} \times 10^2 \times \frac{\pi}{15})$

$\therefore A = \frac{1}{2} \times \frac{\pi}{15} (20^2 - 10^2)$
 $A = 10\pi \text{ cm}^2$ (2)

$f(x) = 2 \cos 2x$ (2)
 $f'(x) = -4 \sin 2x$
 $\therefore f'(\frac{\pi}{4}) = -4 \sin \frac{\pi}{2}$
 $= -4$

Question 2 (12 mks)

a) i) $\frac{d}{dx} (\sin x^2) = \frac{2x \cos x^2}{1}$ (1)

ii) $\frac{d}{dx} (\tan x)^2 = 2 \sec^2 x \cdot \tan x$ (2)

b) $\frac{dy}{dx} = 6 \cos 2x - \sin x$

$y = 3 \sin 2x + \cos x + c$

sub (0,0)
 $0 = 3 \sin 0 + \cos 0 + c$
 $\therefore c = -1$

$y = 3 \sin 2x + \cos x - 1$ (3)

c) $y = 3 \sin x - 4 \cos x$

$\frac{dy}{dx} = 3 \cos x + 4 \sin x$

$\frac{d^2y}{dx^2} = -3 \sin x + 4 \cos x$

LHS = $\frac{d^2y}{dx^2} + y$


$= (-3 \sin x + 4 \cos x) + 3 \sin x - 4 \cos x$
 $= \text{RHS}$ (3)

d) $\int_{\frac{\pi}{3}}^{\frac{2\pi}{3}} \cos 2x \, dx$ (3)

$= \left[\frac{1}{2} \sin 2x \right]_{\frac{\pi}{3}}^{\frac{2\pi}{3}}$
 $= \frac{1}{2} \left[\sin \frac{4\pi}{3} - \sin \frac{2\pi}{3} \right]$
 $= \frac{1}{2} \left[\sin(\pi + \frac{\pi}{3}) - \sin(\pi - \frac{\pi}{3}) \right]$
 $= \frac{1}{2} \left[-\sin \frac{\pi}{3} - \sin \frac{\pi}{3} \right]$
 $= \frac{1}{2} \left[-\frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{2} \right] = -\frac{\sqrt{3}}{2}$

Question 3 (13 mks)

a) $\int_0^2 (1-2x)^3 dx = \left[\frac{(1-2x)^4}{-8} \right]_0^2$
 $= \frac{81}{-8} - \frac{1}{-8}$
 $= -10$ (3)



ii) pt (1, 0) (1)


iii) $A = 2 + \int_0^1 (x^2 - 1) dx + \int_1^3 (x^2 - 1) dx$
 $= 2 + \left[\frac{x^3}{3} - x \right]_0^1 + \left[\frac{x^3}{3} - x \right]_1^3$
 $= 2 + \left(\frac{1}{3} - 1 \right) + \left(6 - \frac{2}{3} \right)$
 $= 2 + \frac{2}{3} + 6\frac{2}{3}$
 $= 9\frac{1}{3} \text{ unit}^2$ (4)

c) $2 \sin^2 x - \sin x = 0$
 $\sin x (2 \sin x - 1) = 0$
 $\sin x = 0 \quad 2 \sin x - 1 = 0$
 $\sin x = \frac{1}{2}$

$\therefore x = 0, \pi, 2\pi, \frac{\pi}{6}, \frac{5\pi}{6}$ (3)

Question 4 (11 mks)

a) i) amp = 4 (1)
period = 4π (1)



ii) $\sin \frac{x}{2} = \frac{1}{4}$
 $4 \sin \frac{x}{2} = 1$
 see $y=1$ above \therefore
2 solutions (1)


b) $\int_1^2 f(x) dx =$
 $= \frac{25}{3} [0 + 1 \cdot 22 + 4 \cdot (67 + 1 \cdot 09) + 2x \cdot 91]$
 $= 0.84$ (2)

c) $V_x = \pi \int_0^3 (\sqrt{x^3})^2 dx$
 $= \pi \int_0^3 x^3 dx$
 $= \pi \left[\frac{x^4}{4} \right]_0^3$
 $= \frac{81\pi}{4} \text{ unit}^3$ (3)

d) $u = x \quad v = \sin 2x$
 $u' = 1 \quad v' = 2 \cos 2x$
 $\therefore \frac{d}{dx} (x \sin 2x) = \underline{\underline{\sin 2x + 2x \cos 2x}}$ (2)

Question 5 (12 mks)

i) i) $y = 2x \quad 3y = x^2$
 $3(2x) = x^2$
 $6x = x^2$
 $0 = x^2 - 6x$
 $0 = x(x-6)$
 $\therefore x = 0 \quad x = 6$ (1)



iii) $A = \int_0^6 (2x - \frac{x^2}{3}) dx$
 $= \left[\frac{2x^2}{2} - \frac{x^3}{9} \right]_0^6$
 $= 36 - \frac{216}{9}$
 $= 12 \text{ unit}^2$ (2)

c) i) $2h + x = 30$
 $2h = 30 - x$
 $h = \frac{30 - x}{2}$
 $\therefore \text{height} = \underline{\underline{\frac{30 - x}{2}}}$ (1)

ii) $A = x \left(\frac{30 - x}{2} \right)$

$A = 15x - \frac{x^2}{2}$ (1)

iii) $\frac{dA}{dx} = 15 - x$ (1)

$\frac{d^2A}{dx^2} = -1$

\therefore st pt $\frac{dA}{dx} = 0 \quad 15 - x = 0$ (3)
 $x = 15$

when $x = 15 \quad \frac{d^2A}{dx^2} < 0 \therefore \text{max}$

c) $\int \tan^2 \theta d\theta$

since $1 + \tan^2 \theta = \sec^2 \theta$

$\int \tan^2 \theta d\theta = \int (\sec^2 \theta - 1) d\theta$
 $= \underline{\underline{\tan \theta - \theta + c}}$ (2)