


Name _____

Teacher _____



Year 12 Mathematics
Assessment Task

Weight: 27%

March 15th 2007

Instructions to students

- Calculators may be used.
- Use black or blue pen only except for graphs.
- Write your name on EVERY page of your solution sheet.
- The phrase 'not attempted' MUST be written on the answer sheet for any question or parts of questions that are not attempted. Insert a blank sheet if an entire question is not attempted.
- Begin each question on a new page.

Time allowed: 1 hour 30 minutes

OUTCOMES ASSESSED	TOPICS COVERED	Result
H1, H2, H4, H5, H6, H9	Probability	/15
	Geometrical Applications of Calculus	/33
	Total	/48

Question 1 Probability


Mark

- (a) A box contains 8 red and 11 green marbles. Shez randomly selects three marbles one at a time and without replacement. What is the probability that she selects *green, red, green* in that order? 1
- (b) In a large school, the student population is 42% male and 58% female. Two students are selected at random to take part in a survey. Find correct to 2 decimal places, the probability that:
- (i) both are female 1
 - (ii) they are different sexes 2
 - (iii) both are of the same sex 2
- (c) Four girls' names and five boys' names are in a hat. Two names are drawn out without replacement. Find the probability that the names of a boy and a girl are drawn out? 2
- (d) In a large barrel there is a mixture of Fundraiser chocolates. White Chocolate Freddos and Strawberry Freddos are in the ratio 2:3. Find the probability that if three chocolates are chosen from the mix, at least one is a Strawberry Freddo? 2
- (e) A coin is biased so that $P(H) = \frac{1}{3}$. How many times must the coin be tossed so that the probability of getting at least one head is more than 90%? 3
- (f) An urn contains W white and B black marbles. Marbles are selected at random, without replacement. 2
- If the probability of selecting 2 white marbles is $\frac{1}{3}$ while the probability of selecting 3 white marbles is $\frac{1}{6}$, find the number of white marbles in the urn.

Question 2 Geometrical Applications of Calculus (start a new page)

Mark

- (a) For the function $y = 2x^3 - 15x^2 + 36x$ determine the values of x for which the function is increasing. 3
- (b) For the curve $y = x^4 - 3x^3 + 9x$ find the value of $\frac{d^2y}{dx^2}$ at the point $(1, 7)$. 2
- (c) Find the equation of the curve $y = f(x)$ if $f''(x) = 2(x-1)$ and the point $(3, -7)$ is a minimum point on the curve. 3
- (d) Find the primitive functions for each of the following
- (i) $3x^4 - 2$ 1
- (ii) $x^{-\frac{1}{3}}$ 2
- (iii) $x^2\sqrt{x}$ 2
- (e) The curve $f(x) = x^3 + 3x^2 - 9x - 1$ is defined in the domain $-4 \leq x \leq 2$. 3
- (i) Find the coordinates of the two stationary points and determine their nature. 4
- (ii) Show that a point of inflexion occurs at $x = -1$. 2
- (iii) Sketch the curve. 3



The sum of the radius R and height h of a cylinder is 60 cm.

- (i) Express h in terms of R . 1
- (ii) Show that the volume of the cylinder is given by $V = \pi(60R^2 - R^3)$. 2
- (iii) Hence find the radius that will give the maximum volume possible. (leave your answer in exact form) 3



Question 2 continues over the page

- (g) The graph of the function Y , defined in the interval $0 \leq x \leq 6$, passes through the points $(2, 10)$, $(3, 15)$ and $(5, 0)$.

The graphs of $\frac{dY}{dX}$ and $\frac{d^2Y}{dX^2}$ are shown below.

Without finding the equation of Y :

- (i) Determine the maximum and minimum points of the graph of Y . 3/3
- (ii) Sketch the graph of Y , labelling the important features. 2/2



QUESTION 1

a) 8r, 11g choose 3

$$P(g, r, g) = \frac{11}{19} \times \frac{8}{18} \times \frac{10}{17} = \frac{440}{2907} \quad (1)$$

b) 420m, 5810f choose 2

I) $P(f, f) = 0.58^2 = 0.3364 \div 0.34$ (2dp) (1) ✓

II) $P(f, m \text{ or } m, f) = 2 \times 0.42 \times 0.58 = 0.4872 \div 0.49$ (2dp) (2)

III) $P(f, f \text{ or } m, m) = 0.58^2 + 0.42^2 = 0.5128 \div 0.51$ (2dp) (2)

c) 4g, 5b choose 2

$$P(b, g \text{ or } g, b) = \frac{5}{9} \times \frac{4}{8} + \frac{4}{9} \times \frac{5}{8} = 2 \times \frac{5}{9} \times \frac{4}{8} = \frac{5}{9} \quad (2) \quad \checkmark$$

d) W:D = 2:3 choose 3

$$P(\text{at least one D}) = 1 - P(\text{no D})$$

$$= 1 - \left(\frac{2}{5}\right)^3 \quad \checkmark$$

$$= 1 - \frac{8}{125}$$

$$= \frac{117}{125} (= 0.936) \quad \checkmark \quad (2)$$

e) $P(H) = \frac{1}{3} \Rightarrow P(T) = \frac{2}{3}$

$$P(\text{at least one H}) = 1 - P(\text{no H})$$

$$= 1 - P(\text{all T})$$

$$= 1 - \left(\frac{2}{3}\right)^n \quad \checkmark$$

is $0.90 < 1 - \left(\frac{2}{3}\right)^n \quad \checkmark$

$$-0.1 < -\left(\frac{2}{3}\right)^n$$

$$0.1 > \left(\frac{2}{3}\right)^n$$

$$\frac{\ln 0.1}{\ln \frac{2}{3}} < n$$

$$n > 5.7$$

$\therefore n = 6 \quad \checkmark$

(3)

f) $P(2w) = \frac{1}{3}, P(3w) = \frac{1}{6}$

let no of marbles be n .

now $P(2w) = \frac{1}{3} = \frac{w}{n} \times \frac{w-1}{n-1}$
 $= \frac{w^2 - w}{n^2 - n}$

$$n^2 - n = 3w^2 - 3w \quad (1)$$

now $P(3w) = P(2w) \times \frac{w-2}{n-2}$

is $\frac{1}{6} = \frac{1}{3} \times \frac{w-2}{n-2}$

$$\frac{1}{2} = \frac{w-2}{n-2}$$

$$n-2 = 2w-4$$

$$n = 2w-2 \quad (2) \quad \checkmark$$

into (1).

$$(2w-2)^2 - (2w-2) = 3w^2 - 3w$$

$$4w^2 - 8w + 4 - 2w + 2 = 3w^2 - 3w$$

$$w^2 - 7w + 6 = 0$$

$$(w-6)(w-1) = 0$$

$$\therefore w = 1, 6$$

but $w \neq 1 \therefore w = 6$.

The minimum no of white marble is 6. ✓ (2)

\therefore The no of white marbles is $6n, n \in \mathbb{Z}^+$

Question 2.

$$y = 2x^3 - 15x^2 + 36x$$

$$y' = 6x^2 - 30x + 36$$

increasing when $y' > 0$

$$6x^2 - 30x + 36 > 0$$

$$x^2 - 5x + 6 > 0$$

$$(x-3)(x-2) > 0$$

$$\therefore x < 2, x > 3$$

$$\therefore f'(x) = x^2 - 2x - 3$$

$$f(x) = \frac{x^3}{3} - x^2 - 3x + C_2$$

When $x = 3, y = -7$

$$\therefore -7 = 9 - 9 - 9 + C_2$$

$$2 = C_2$$

$$\therefore f(x) = \frac{x^3}{3} - x^2 - 3x + 2$$

b) $y = x^4 - 3x^3 + 9x$

$$\frac{dy}{dx} = 4x^3 - 9x^2 + 9$$

$$\frac{d^2y}{dx^2} = 12x^2 - 18x$$

at $x = 1$

$$\frac{d^2y}{dx^2} = 12 - 18$$

$$= -6$$

c) $f''(x) = 2x - 2$

$$f'(x) = x^2 - 2x + C_1$$

when $f'(x) = 0, x = 3$

$$0 = 9 - 6 + C_1$$

$$-3 = C_1$$

d) i) $\int 3x^4 - 2 dx$

$$= \frac{3x^5}{5} - 2x + C_1$$

ii) $\int x^{-\frac{1}{3}} dx$

$$= \frac{3}{2} x^{\frac{2}{3}} + C_2$$

iii) $\int x^2 \sqrt{x} dx$

$$= \int x^{\frac{5}{2}} dx$$

$$= \frac{2}{7} x^{\frac{7}{2}} + C_3$$

2e)

$$f(x) = x^3 + 3x^2 - 9x - 1$$

$$-4 \leq x \leq 2$$

(iii) $x = -4, y = 19 \quad (-4, 19)$

$$x = 2, y = 1 \quad (2, 1)$$

i) for st pts $f'(x) = 0$

$$x = 0, y = -1 \quad (0, -1)$$

$$f'(x) = 3x^2 + 6x - 9$$

$$0 = 3x^2 + 6x - 9$$

$$0 = x^2 + 2x - 3$$

$$0 = (x+3)(x-1)$$

$$x = -3 \text{ or } 1$$

x	-4	-3	-2
$f'(x)$	+15	0	-9

\therefore max at $(-3, 26)$

\therefore min at $(1, -6)$

x	0	1	2
$f'(x)$	-9	0	+15

\therefore min at $(1, -6)$

ii) For POT $f''(x) = 0$

$$f''(x) = 6x + 6$$

If $f''(x) = 0$


$$x = -1$$

\therefore possible inf at $(-1, 10)$

x	-2	-1	0
$f''(x)$	-6	0	+6

Change of concavity

\therefore inf at $(-1, 10)$



f) i) $R + h = 60$

$$h = 60 - R$$

ii) $V = Ah$

$$= \pi R^2 (60 - R)$$

$$= \pi (60R^2 - R^3)$$

iii) $V = 60\pi R^2 - \pi R^3$

$$V' = 120\pi R - 3\pi R^2$$

for max/min $V' = 0$

$$0 = 3\pi R(40 - R)$$

$$\therefore R = 40, R \neq 0$$

$$V'' = 120\pi - 6\pi R$$

if $R = 40$

$$V'' < 0$$

\therefore max volume

when $R = 40$

Q2 ctd

g) $0 \leq x \leq 6$

$(2, 10)$ $(3, 15)$ $(5, 0)$

i) $\min(2, 10)$ |

$\max(3, 15)$ |

$\min(5, 0)$ |

$x \cdot 2 \cdot 3 \cdot 5 \cdot$

$\frac{dy}{dx} - 0 + 0 - 0 +$

ii)

