



2011  
HIGHER SCHOOL  
CERTIFICATE  
ASSESSMENT 3

# Mathematics

## General Instructions

- Working Time - 45 mins.
- Write using a blue or black pen.
- Approved calculators may be used.
- All necessary working should be shown for every question.
- Begin each question on a fresh sheet of paper.

Total marks (30)


- Attempt Questions 1-3.
- All questions are of equal value.

*Remember to use the standard integrals sheet*

Question 1	(10 Marks)	Start a new page	Marks
<del>1</del>	Differentiate with respect to $x$		
<del>1</del>	$xe^x$		2
<del>1</del>	$\frac{x+2}{\tan x}$		2
<del>1</del>	$(\log_e x + 1)^3$		2
<del>1</del>	Evaluate		
<del>1</del>	$\int \frac{10x}{x^2 + 1} dx$		2
<del>1</del>	$\int \sec^2 3x dx$		2
<b>Question 2</b>	<b>(10 Marks)</b>	<b>Start a new page</b>	<b>Marks</b>
<del>1</del>	The curve $y = e^{2x}$ is rotated about the $x$ -axis.		
	Show that the volume of the solid of revolution formed between $x = 0$ and $x = 3$ is given by $\frac{\pi}{4}(e^{12} - 1)u^3$		3

**Question 2** continued

- (b) The diagram shows a circle with centre  $O$  and radius  $7\text{cm}$ .  
The length of the arc  $LM$  is  $10\text{cm}$ .



- (i) Find  $\angle LOM$  1
- (ii) Hence find the area of the minor segment bounded by  $LM$  correct to 1 decimal place 2

- (c) The rate at which people flow into a train station during afternoon peak hour is given by:

$$\frac{dP}{dt} = 60 - 16t \quad \text{for } t \geq 0$$

where  $P$  is the number of people and  $t$  is the time in measured in hours.

- (i) Find the initial rate at which people flow into the station 1
- (ii) Find the number of people in the station as a function of  $t$ . 1
- (iii) Initially, there are 240 people in the station.  
Find out how many people remain in the station after 5 hours 2

**Question 3** (10 Marks) Start a new page Marks

- (a) A particle moves in a straight line and at any time  $t$  seconds  $t \geq 0$ , its displacement  $x(t)$  from 0 is measured in metres

Its velocity in metres per second is given by:

$$x(t) = 1 - \sin(2t) \quad \text{for } 0 \leq t \leq 2\pi$$

- (i) Find when the particle is first at rest 2
- (ii) Find an expression for the acceleration 1
- (iii) The particle is initially  $1.5\text{m}$  from  $O$ .  
Find an expression for the displacement  $x(t)$ . 2

- (b) Evaluate  $\frac{d}{dx} \sin(x^{-3})$  2

- (c) Current  $i$  is measured in Amps and time is measured in seconds.

A current  $i_0$  is established in an electrical circuit. After the source of the current has been removed, the current in the circuit decays according to the equation  $\frac{di}{dt} = -ki$

- (i) Show that  $i = i_0 e^{-kt}$  satisfies the differential equation  $\frac{di}{dt} = -ki$ . 1
- (ii) The current in the circuit decays to 36.8% of the original current in a 0.1 of a second. Find  $k$  to 1 decimal place. 2

(Q1)

Q1. a) i)  $y = xe^x$

$u = x$      $v = e^x$   
 $u' = 1$      $v' = e^x$

(10)

$y' = vu' + uv'$   
 $y' = e^x + xe^x$

(2)

ii)  $y = \frac{x+2}{\tan x}$

let  $u = x+2$      $v = \tan x$   
 $u' = 1$      $v' = \sec^2 x$

$y' = vu' - uv'$

$\therefore y' = \frac{\sqrt{2} \tan x - (x+2)(\sec^2 x)}{(\tan x)^2}$

(2)

iii)  $y = (\log_e x + 1)^3$

$\therefore y' = 3(\log_e x + 1)^2 \cdot \left(\frac{1}{x}\right)$

(2)

b) i)  $\int \frac{10x}{x^2+1} dx$

$u = x^2+1$   
 $u' = 2x$

$= 5 \int \frac{2x}{x^2+1} dx$

$= 5 \ln(x^2+1) + c$

(2)

ii)  $\int \sec^2 3x dx$      $a = 3$


$= \frac{1}{3} \tan 3x + c$

(2)

(Q2)

a)  $y = e^{2x}$

$V = \pi \int_0^3 y^2 dx$



(10)

$y = e^{2x}$   
 $\therefore y^2 = e^{4x}$

$\therefore V = \pi \int_0^3 e^{4x} dx$

$= \pi \left[ \frac{1}{4} e^{4x} \right]_0^3$

$= \frac{\pi}{4} \left[ e^{4x} \right]_0^3$

$= \frac{\pi}{4} [e^{12} - e^0]$

$= \frac{\pi}{4} [e^{12} - 1] u^3$  as req.

(3)

$$b) i) l = r\theta$$

$$10 = 7\theta$$

$$\therefore \theta = \frac{10}{7} \text{ radians}$$

✓ (1)

minor segment:

$$ii) A = \frac{1}{2} r^2 (\theta - \sin \theta) \quad \text{1 dp}$$

$$A = \frac{1}{2} r^2 \left( \frac{10}{7} - \sin \frac{10}{7} \right)$$

$$= \frac{1}{2} (7)^2 \left( \frac{10}{7} - \sin \left( \frac{10}{7} \right) \right) \quad (\text{radians mode})$$

$$\therefore A = 10.747$$

$$= 10.7 \text{ cm}^2 \quad (1 \text{ dp})$$

✓ (2)

$$c) \frac{dP}{dt} = 60 - 16t \quad t \geq 0$$

i) initial @  $t=0$

$$\frac{dP}{dt} = 60 - 16(0)$$

$$= 60 \text{ people/hr.} \quad \checkmark (1)$$

$$ii) \int \frac{dP}{dt} = 60 - 16t \quad dt$$

$$\therefore P = 60t - \frac{16t^2}{2} + C$$

$$\therefore P = 60t - 8t^2 + C \quad \checkmark (1)$$

iii)  $t=0, P=240$ .

$$~~P = 60t - 8t^2 + C~~$$

$$P = 60t - 8t^2 + C$$

$$240 = 60(0) - 8(0) + C$$

$$\therefore C = 240 \quad \checkmark$$

$$\therefore P = 60t - 8t^2 + 240$$

( $P=?$   $t=5$ )

$$P = 60(5) - 8(5)^2 + 240$$

$$\therefore P = \boxed{340 \text{ ppl}} \text{ remain at station after 5 hrs.} \quad \checkmark (2)$$

remainings

BRNO

3

a)  $t \geq 0$  meters

m/sec.

10

$$x = 1 - \sin(2t)$$

$$0 \leq t \leq 2\pi$$

i) rest when  $v=0$

$$v = 1 - \sin(2t)$$

$$1 - \sin 2t = 0$$

$$\sin 2t = 1$$

$$2t = \frac{\pi}{2}$$

$$\therefore t = \frac{\pi}{4} \text{ sec. first @ rest.}$$

ii) find  $\ddot{x}$

differentiate

$$x = 1 - \sin(2t)$$

$$\dot{x} = -2 \cos(2t)$$

SC-S-C

iii)  $t=0, x = 1.5 \text{ cm}$

$$\int \dot{x} = 1 - \sin(2t) dt$$

$$x = t + \frac{1}{2} \cos(2t) + c$$

(@  $t=0, x=0.015$ )

$$0.015 = 0 + \frac{1}{2} \cos(0) + c$$

$$\therefore 0.015 = \frac{1}{2}(1) + c$$

$$c = -0.485$$

$$\therefore x(t) = t + \frac{1}{2} \cos(2t) - 0.485$$

m/s.

PTO

SC-S-C

$$a) \text{iii) } \int \dot{x} = 1 - \sin 2t dt$$

$$x = t + \frac{1}{2} \cos 2t + c$$

$$x = 1.5 \text{ m, } t=0$$

$$1.5 = 0 + \frac{1}{2} \cos 2(0) + c$$

$$1.5 = 0 + \frac{1}{2} + c$$

$$c = 1.5 - 0.5$$

$$c = 1$$

$$\therefore x(t) = t + \frac{1}{2} \cos 2t + 1 \text{ for m/s.}$$

2

5c - s-c

b)  $y = \sin(x^{-3})$

$y' = -3x^{-4} \cos(x^{-3})$  let  $u = x^{-3}$

$u' = -3x^{-4}$

$\therefore \frac{dy}{dx} = \frac{-3}{x^4} \cos\left(\frac{1}{x^3}\right)$

$u = x^{-3}$   
 $u' = -3x^{-4}$

(2)

e) i)  $i = i_0 e^{-kt}$

$\frac{di}{dt} = -K(i_0 e^{-kt})$

since  $i = i_0 e^{-kt}$

$\therefore \frac{di}{dt} = -K(i)$

$\frac{di}{dt} = -ki$  as req.

(1)

ii) decays to 36.8%

hence  $I =$

$i = i_0 e^{-kt}$

$i = \text{current (amps)}$

because it decays to 36.8% of original

$36.8\% = 0.368$

$\therefore i = 0.368 i_0$

$i = i_0 e^{-kt}$

subbing in

$0.368 i_0 = i_0 e^{-kt}$

$e^{-kt} = 0.368$   $t = 0.1 \text{ sec.}$

$e^{-k(0.1)} = 0.368$

$\therefore k = \frac{\ln(0.368)}{-0.1}$

(2)

hence  $k = 9.9967$

$k = 10.0$  (1 dp)