

Name. _____

Teacher. _____



MORIAH COLLEGE

Year 11

MATHEMATICS PRELIM

5th September 2014

Time Allowed: 2 hours


Examiners: P. Brown, L Bornstein

Instructions:

- Answer every question.
- Show all necessary working. Draw clear, well labelled diagram.
- There are 10 multiple choice questions each worth 1 mark.
- There are 4 long questions each worth 15 marks.
- Marks may be deducted for careless or untidy work.
- Board approved calculators may be used.

1.	Simplify the expression $\frac{x^2 + 4x + 4}{x + 2}$. (A) $x + 2$ (B) $\frac{x^2 + 2x + 2}{x}$ (C) $3x + 4$ (D) 1			
2.	What is the angle of inclination that the line $3x - 5y - 7 = 0$ makes with the positive direction of the x-axis? (A) $149^\circ 2'$ (B) $59^\circ 2'$ (C) $120^\circ 28'$ (D) $30^\circ 58'$			
3.	Make r the subject of the formula: $a = rs + 1$ (A) $r = a - s$ (B) $r = \frac{a - 1}{s}$ (C) $r = \frac{a - s}{s}$ (D) $r = \frac{a + s}{s}$			
4.	Find the exact value of $\cos 210^\circ$ (A) $-\frac{\sqrt{3}}{2}$ (B) $-\frac{1}{2}$ (C) $\frac{1}{2}$ (D) $-\frac{\sqrt{3}}{2}$			
5.	What is the value of $f(-1)$ if $f(x) = x^3 - 4x$? (A) -3 (B) -5 (C) 3 (D) 5			
6.	For the arithmetic series 3, 6, 9, ... which expression could be used to evaluate S_{15} ? (A) $3 + 14(3)$ X (B) $15[3 + 14(3)]$ (C) $\frac{15}{2}[3 + 14(3)]$ X (D) $\frac{15}{2}(6 + 14(3))$			
7.	The gradient of the normal to the curve $f(x) = 3x^3 - 4x + 2$ at the point (-1, 3) is: (A) 5 (B) -5 (C) $-\frac{1}{5}$ (D) $-\frac{1}{3}$			

8. Which of the equations given below describes the circle centre O, shown?



- (A) $x^2 - 4x + y^2 - 2y = 21$ (B) $(x - 2)^2 + (y - 1)^2 = 16$
 (C) $x^2 + y^2 = 16$ (D) $(x + 2)^2 + (y + 1)^2 = 16$

9. If $f(x) = \frac{2}{x}$ and $g(x) = 3 - x$; find the values of x if $f(x) = g(x)$,
 (A) $x = 1$ and $x = 2$ (B) $x = -1$ and $x = 2$
 (C) $x = -1$ and $x = -2$ (D) $x = 1$ and $x = -2$

10. What is the solution to the equation $\sin(\frac{\theta}{2} + 30^\circ) = \cos \theta$

- (A) $\theta = 40^\circ$ (B) $\theta = 60^\circ$ (C) $\theta = 80^\circ$ (D) $\theta = 100^\circ$

Question 11 (start in a new booklet) (15 marks)

a. Evaluate $\frac{3.7^2}{\cos^2 54^\circ 18'}$ correct to three significant figures.

b. Fully simplify $\frac{1}{2\sqrt{3}-4}$ by rationalising the denominator

c. Express the following in simplest terms

$$\frac{x^2 - 3x - 4}{x^2 - 1} \times \frac{x^2 - x}{x^2 - 2x - 8}$$

d. Solve $2\cos x + 1 = 0$ for $0^\circ \leq x \leq 360^\circ$


e. Solve the following inequality and sketch the solution on the number line.

$$|3x - 2| \geq 4$$

f. Prove that: $\sin^2 \theta \cdot \cos^2 \theta + \sin^4 \theta = \sin^2 \theta$

Question 12 (start in a new booklet) (15 marks)


a.



$A(-2, 2)$, $B(-1, -5)$ and $C(6, -6)$ are vertices of a triangle and P lies on the line AC .

- i) Show that the midpoint P of AC has coordinates $(2, -2)$. 1
- ii) Find the gradient of BP . 1
- iii) Show that BP is perpendicular to AC . 1
- iv) Show that the line BP has equation $x - y - 4 = 0$. 2
- v) Find the coordinates of D if P is the midpoint of BD . 2
- vi) Find the perpendicular distance from A to BP . 2

b.



A , B , and C are markers in an orienteering course. $AC = 4 \text{ km}$ and $BC = 5 \text{ km}$. The bearing of C from B is 40°T . The bearing of B from A is 260°T

- i) Copy the diagram into your books and, clearly mark all information onto your diagram. Hence show that $\angle CAB = 40^\circ$ 2
- ii) Use the Sine Rule to show that $\angle CAB = 53^\circ$ to the nearest degree. 2
- iii) Hence find the bearing of C from A . 2

Question 13 (start in a new booklet) (15 marks)

a. Find: $\lim_{x \rightarrow 1} \frac{x^2 + 3x - 4}{x - 1}$

1

- b. Differentiate each of the following expressions with respect to x.
(fully simplify all answers)

i) $3x^3 + 3x^2 - 2x + 1$

1

ii) $(2x - 1)^6$

1

iii) $\frac{x+1}{3x-2}$

2

iv) $\frac{2}{\sqrt{x}} + 2\sqrt{x}$

2

(give solution without negative fractional indices)

- c. Consider the curve given by $y = x^3 - 3x^2 - 9x + 20$.

- i) Find the coordinates of the two stationary points and determine their nature.

4

- ii) Find the point of inflexion.

2

- iii) Sketch the curve, showing any other relevant features if $-2 \leq x \leq 5$

2

Question 14 (start in a new booklet) (15 marks)

- a. Solve for θ if $2\cos^2\theta - 3\sin\theta = 0$ in the domain $0^\circ \leq \theta \leq 360^\circ$

3

- b. Given $y = \sqrt{1+x^2}$, show that $\frac{dy}{dx} = \frac{x}{y}$

2

- c. The tangent to $y = ax^3 + bx - 2$ at $(-1, 2)$ is horizontal.
Find a and b .

3

- d. A function $f(x)$ is defined as

$$f(x) = \begin{cases} 5x - 4 & \text{for } x \geq 0 \\ px + q & \text{for } x < 0 \end{cases}$$

Find p and q , so that $f(x)$ is an even function.

2

- e. Draw a sketch of the function $y = f(x)$ given that $f(x)$ has the following properties:

$f(x)$ is defined over the domain $-1 \leq x \leq 1$

$f(x)$ is negative for $x < 0$ and is positive for $x > 0$.

$f(x)$ is increasing everywhere.

$f(x)$ is an odd function

2

- f. Write 3 terms of an arithmetic sequence whose product is 315 and whose sum is 21.

3

END OF TEST

Section I

10 marks

Attempt Questions 1-10

Allow about 15 minutes for this section

Select the alternative A, B, C or D that best answers the question and indicate your choice with a cross (X) in the appropriate space on the grid below.

	A	B	C	D
1	X			
2				X
3	X			
4			X	
5		X		
6			X	
7		X		
8	X			
9	X			
10	X			

Y11 24 2014
prelim.

#1a. $40 \cdot 20317 \checkmark$

$\therefore 40.2 (\text{to } 3 \text{ s.f.}) \checkmark$

b. $\frac{1}{(2\sqrt{3}-4)} \times \frac{2\sqrt{3}+4}{2\sqrt{3}+4} = \frac{2\sqrt{3}+4}{12-16} = \frac{2\sqrt{3}+4}{-4}$
 $= \frac{\sqrt{3}+2}{-2} \checkmark$

c. $\frac{(x-4)(x+1)\cancel{x}(x-1)\cancel{x}}{(x-1)(x+1)(x-4)(x+2)} = \frac{x}{x+2} \checkmark$

d. $\cos \alpha = -\frac{1}{2}$
 $\text{reflex } 60^\circ \checkmark$
 2nd | 3rd.
 $\alpha = 120^\circ \quad \alpha = 240^\circ \checkmark$

e. Consider $|3x-2| = 4$.

either $3x-2 = 4$ or $-(3x-2) = 4$

$3x = 6 \quad 3x-2 = -4$

$x = 2 \quad 3x = -2$

$x = -\frac{2}{3}$ $x=0$ $x=2$. $x=3$

$x \leq -\frac{2}{3} \checkmark \quad x \geq 2 \checkmark$

f. LHS:

$$\begin{aligned} & \sin^2 \theta \cos^2 \theta + \sin^4 \theta \\ &= \sin^2 \theta (\cos^2 \theta + \sin^2 \theta) \\ &= \sin^2 \theta (1) \\ &= \sin^2 \theta \end{aligned}$$

RHS:

$\sin^2 \theta$

∴

LHS = RHS

$$\#13a \lim_{x \rightarrow 1} \frac{(x+4)(x+1)}{(x-1)} = 5\sqrt{1}$$

$$b.i) y = 3x^3 + 3x^2 - 2x + 1 \\ y' = 9x^2 + 6x - 2.$$

$$ii) y = (2x-1)^6 \\ y' = 6(2x-1)^5 \cdot 2 \\ y' = 12(2x-1)^5.$$

$$iii) y = \frac{x+1}{3x-2}$$

$$u = x+1 \quad v = 3x-2 \\ u' = 1 \quad v' = 3.$$

$$y' = \frac{(3x-2)(1) - 3(x+1)}{(3x-2)^2}$$

$$y' = \frac{3x-2 - 3x-3}{(3x-2)^2} \\ y' = \frac{-5}{(3x-2)^2}$$

$$iv) y = 2x^{-\frac{1}{2}} + 2x^{\frac{1}{2}}$$


$$y' = -x^{-\frac{3}{2}} + 1x^{-\frac{1}{2}} \\ y' = \frac{-1}{\sqrt{x^3}} + \frac{1}{\sqrt{x}}$$

$$c. y = x^3 - 3x^2 - 9x + 20$$

$$P.O.: y' = 0 \\ y' = 3x^2 - 6x - 9 = 0 \\ 3(x^2 - 2x - 3) = 0 \\ 3(x-3)(x+1) = 0 \\ x=3 \quad x=-1 \\ y = -7 \quad y = 25 \\ (-3, -7) \quad (-1, 25) \\ \text{min.} \quad \text{max}$$

Nature:

x	-2	-1	0	3	4
y'	15	0	-9	0	15



$$P.O.I.: y'' = 0$$

$$y'' = 6x - 6 = 0 \\ x = 1$$

$$y = 9 \\ \text{possible P.O.I. } (1, 9)$$

x	0	1	2
y''	-6	0	6

\therefore concavity changes.

$$x = -2 \quad x = 5 \\ y = 18 \quad y = 25$$

$$\#14a. 2\cos^2\theta - 3\sin\theta = 0.$$

$$2(1-\sin^2\theta) - 3\sin\theta = 0.$$

$$2 - 2\sin^2\theta - 3\sin\theta = 0.$$

$$\sin\theta = 2\sin^2\theta + 3\sin\theta - 2 \quad \checkmark$$

$$\sin\theta = (2\sin\theta - 1)(\sin\theta + 2)$$

$$\sin\theta = \frac{1}{2}$$

$$\sin\theta = -2$$

$$\text{red } \lambda = 30^\circ$$

$$\text{no soln.} \quad \checkmark$$

3.

$$1st \quad | \quad 2nd \quad \checkmark$$

$$\theta = 30^\circ \quad | \quad \theta = 150^\circ$$

\rightarrow

$$b. \quad y = ax^3 + bx - 2$$

$$(-1, 2) : 2 = a(-1)^3 + b(-1) - 2$$

$$2 = -a - b - 2$$

$$a + b = -4 \quad \text{--- (1)} \quad \checkmark$$

$$y' = 3ax^2 + b = 0$$

at $x = -1$.

$$3a(-1)^2 + b = 0.$$

$$3a + b = 0$$

$$b = -3a \quad \text{--- (2)} \quad \checkmark$$

Subst (2) in (1).

$$a - 3a = -4$$


$$-2a = -4$$

$$a = 2.$$

$$b = -6.$$

3.

$$c. \quad F(x) = \begin{cases} 5x - 4 & x \geq 0 \\ px + q & x < 0. \end{cases}$$



$$d. \quad y = (1+x^2)^{\frac{1}{2}}$$


$$\frac{dy}{dx} = \frac{1}{2}(1+x^2)^{-\frac{1}{2}} \cdot 2x \quad \checkmark$$

$$= x \quad \sqrt{1+x^2} \quad 2.$$

$$= \frac{x}{\sqrt{1+x^2}} \quad \checkmark$$

$$\underline{y}.$$

e.



3.

f) terms: $a-d, a, a+d$:

$$a(a-d)(a+d) = 315$$

$$7(7-d)(7+d) = 315$$

$$49 - d^2 = 45$$

$$49 - 45 = d^2$$

$$4 = d^2$$

$$d = \pm 2. \quad \checkmark$$

$$7-2, 7, 7+2$$

$$5, 7, 9. \quad \checkmark$$

3.

45 mins

#12 i) $P = \text{midpt}(AC) = \left(\frac{-2+6}{2}, \frac{2-6}{2} \right) = (2, -2) \checkmark$

ii) $m(BP) = \frac{-2-5}{2-1} = \frac{3}{1} = 3 \checkmark$

iii) $m(AC) = \frac{-6-2}{6-2} = \frac{-8}{4} = -2$

$\text{scc } 1 \times -1 = -1$
 $BP \perp AC \checkmark$

iv) eq. of BP:

$$y - 5 = 1(x - 1) \checkmark$$

$$y + 5 = x + 1 \quad 2.$$

$$0 = x - y - 4 \checkmark$$

v) $(x_1 - x_2) = \text{midpt } BD$

$$(x_1 - x_2) = \left(\frac{-1+x}{2}, \frac{-5+y}{2} \right)$$


$$\begin{array}{l|l} -1+x = 2 & -5+y = -2 \\ -1+2 = 4 & -5+y = -4 \\ x = 5 \checkmark & y = 1 \checkmark \end{array}$$

$$D(5, 1) \quad 2.$$

vii) A(-2, 2) BP: $x - y - 4 = 0$.

$$d = \sqrt{|-2 - 2 - 4|} \quad 2$$

$$= \sqrt{\frac{-8}{2}} = \frac{8}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{8\sqrt{2}}{2} = 4\sqrt{2} \checkmark$$



$$\sin \theta = \frac{\sin 40^\circ}{5} \checkmark$$

$$\sin \theta = \frac{5 \cdot \sin 40^\circ}{4}$$

$$\sin \theta = 0.803$$

$$\theta = \sin^{-1}(0.803)$$

$$\theta = 53^\circ \checkmark$$

6.

: bearing is $313^\circ T \checkmark$ or $N 47^\circ W$.