



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS

April 2014

Assessment Task 2
Year 12

Mathematics Extension 1

General Instructions

- Reading Time – 5 Minutes
- Working time – 90 Minutes
- Write using black or blue pen. Pencil may be used for diagrams.
- Board approved calculators maybe used.
- Marks may **NOT** be awarded for messy or badly arranged work.
- All necessary working should be shown in every question if full marks are to be awarded.
- Answer in simplest exact form unless otherwise instructed.

Total Marks – 80

- Attempt sections A – D.
- Start each **NEW** section in a separate answer booklet.
- Hand in your answers in 4 separate bundles:
 - Section A
 - Section B
 - Section C
 - Section D

Examiner: *J. Chen*

START A NEW ANSWER BOOKLET

SECTION A [20 marks]

1. If $f(x) = x + 5$, find the inverse function $f^{-1}(x)$. [1]
2. Find the Cartesian equation whose parametric equations are [1]
- $$x = 8t, \quad y = 4t^2$$
3. If α, β, γ are the roots of the equation $x^3 + x^2 - 2x - 1 = 0$. Find the value of
- (i) $\alpha + \beta + \gamma$ [1]
- (ii) $\alpha\beta\gamma$ [1]
- (iii) $\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2$ [1]
4. Let $y = e^x$. Find $\frac{dy}{dx}$ when $x = 1$. [2]
5. Find [2]
- $$\frac{d}{dx} \cos^{-1} 2x$$
6. There are 40 Students in a class. 38 of them are boys. If 5 students are randomly selected, find the number of ways that at least 1 girl is selected. [2]
7. Let $P(x)$ be a polynomial. When $P(x)$ is divided by $x - 2$, the quotient is $6x^2 + 5x + 9$. It is given that $P(2) = 20$.
- (i) Find $P\left(\frac{2}{3}\right)$. [2]
- (ii) Solve $P(x) = 0$ [3]
8. Peter invites 8 friends to join his birthday party. There are 5 boys and 4 girls at the party including Peter. In a game, boys and girls sit in a row. What is the probability that the boys and girls sit alternatively? [2]
9. Find [2]
- $$\int \frac{2}{\sqrt{16 - x^2}} dx$$

End of Section A

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate

START A NEW ANSWER BOOKLET

SECTION B [20 marks]

1. Let $P(x) = x^3 + x^2 + mx + n$. If $P(1) = 0$ and the remainder of $P(x)$ divided by $x - 2$ is 11, then factorise $P(x)$.

Marks
[3]

2. Consider the function $f(x) = \sin^{-1}(x - 2)$, evaluate $f\left(\frac{3}{2}\right)$.

[2]

3. There are 24 boys and 4 girls in a class. From the class, 5 students are randomly selected to form the class committee.

(i) Find the probability that the class committee consists of boys only.

[1]

(ii) Find the probability that the class committee consists of at least 2 boys and 1 girl.

[3]

4. Show that if $t = \tan x$, then

[2]

$$\frac{dx}{dt} = \frac{1}{1+t^2}$$

5. Find

(i)

$$\int \frac{1}{\sqrt{25-9x^2}} dx$$

[2]

(ii)

$$\int \frac{x+1}{x^2+1} dx$$

[2]

6. Find

[2]

$$\int (e^x + e^{-x})^2 dx$$

7. Find the volume of the solid generated when the region bounded by the curve $y = e^x$ and the x -axis in the interval $0 \leq x \leq 2$ is rotated about the x -axis.

[3]

End of Section B

START A NEW ANSWER BOOKLET

SECTION C [20 marks]


1. Find

Marks
[2]

$$\int_0^{\frac{\sqrt{3}}{2}} \frac{dx}{\sqrt{3-4x^2}}$$

2. In the diagram below, the line $L: y = \frac{1}{4 \ln 2} x + \left(2 - \frac{1}{\ln 2}\right)$ touches the curve $C: y = \log_2 x$ at the point P. Find the coordinates of P.

[3]



3. Prove, by mathematical induction, that $9^n - 1$ is divisible by 8 for all positive integers n .

[3]

4. The slope at any point (x, y) of a curve is given by $\frac{dy}{dx} = 2e^{2x} - 3e^x$. The y -intercept of the curve is 0. Find the area bounded by the curve and the x -axis.

[4]

Section C continues on next page

5. (i) Find $\frac{d}{dx}(x^2 \ln x)$ [2]

(ii) Hence, find [3]

$$\int x \ln x \cdot dx$$

6. Sketch $y = 3 \cos^{-1} 2x$. State the domain and range. [3]

End of Section C

START A NEW ANSWER BOOKLET

SECTION D [20 marks]

Marks

1. $P(2ap, ap^2)$ and $Q(2aq, aq^2)$ are two points on the parabola $x^2 = 4ay$. The tangents at P and Q meet at the point T .

(i) Show that the equation of the tangent at P is $y = px - ap^2$ and similarly, write down the equation of the tangent at Q . [2]

(ii) Find the coordinates of T . [2]

(iii) Let M be the midpoint of PQ , find the coordinates of M . [1]

(iv) The point R is the midpoint of MT , show that R has coordinates [1]

$$\left[a(p+q), \frac{a(p+q)^2}{4} \right]$$

(v) Find the Cartesian equation of the locus of R and describe the locus in geometrical terms. [2]

2. Consider the following letters [3]

A, A, B, B, C, C

how many different arrangements are possible if no two identical letters are next to one another?

3. (i) Show that [2]

$$a^n + b^n = (a+b)(a^{n-1} + b^{n-1}) - ab(a^{n-2} + b^{n-2}),$$

where a and b are real numbers.

(ii) Let $x + \frac{1}{x} = 2 \cos \theta$, where $\theta \neq \pi$ and $x \neq 0$. Suppose [3]

$$x^k + \frac{1}{x^k} = 2 \cos k\theta \text{ and } x^{k+1} + \frac{1}{x^{k+1}} = 2 \cos(k+1)\theta \text{ for a positive integer } k.$$

$$\text{Show that } x^{k+2} + \frac{1}{x^{k+2}} = 2 \cos(k+2)\theta.$$

(iii) It is given that $x^n + \frac{1}{x^n} = 2 \cos n\theta$. Prove, by mathematical induction, that [4]

$$(x + x^3 + x^5 + \dots + x^{2n-1}) + \left(\frac{1}{x} + \frac{1}{x^3} + \frac{1}{x^5} + \dots + \frac{1}{x^{2n-1}} \right) = \frac{\sin 2n\theta}{\sin \theta}$$

for all positive integers n , where $\theta \neq k\pi$.

End of Section D
End of Exam

Section A

1. $f(x) = x + 5^{-x}$

ie $y = x + 5^{-x}$

For $f^{-1}(x)$:

$x = y + 5^{-y}$

ie $y = x - 5^{-y}$

So $f^{-1}(x) = x - 5^{-x}$ [1]

2. $x = 8t$; $y = 4t^2$

$t = \frac{x}{8}$; $y = 4\left(\frac{x}{8}\right)^2$

$= \frac{4x^2}{64}$

$\therefore y = \frac{x^2}{16}$ [1]

3. $x^3 + x^2 - 2x - 1 = 0$

(i) $\alpha + \beta + \gamma = -1$ [1]

(ii) $\alpha\beta\gamma = 1$ [1]

(iii) $\alpha^2\beta\gamma + \alpha\beta^2\gamma + \alpha\beta\gamma^2$
 $= \alpha\beta\gamma(\alpha + \beta + \gamma)$

$= +1 \times -1$

$= -1$ [1]

4. $y = e^{2x}$

$\frac{dy}{dx} = e^{2x}$

When $x=1$

$\frac{dy}{dx} = e^2$
 $= e$ [2]

5. $\frac{d}{dx} \cos^{-1} 2x$

$= \frac{-1}{\sqrt{1-(2x)^2}} \times 2$

$= \frac{-2}{\sqrt{1-4x^2}}$ [2]

6. No. of ways of choosing 5 including at least one girl is the total no. of ways of choosing 5, minus the no. of ways of choosing 5 boys.

$40C_5 - 38C_5$
 $= 156066$

[2]

7. Division transformation: [2]

$P(x) = (x-2)(6x^2 + 5x + 9) + 20$

(i) $P\left(\frac{2}{3}\right) = 0$ [2]

(ii) $P(x) = 6x^3 - 7x^2 - x + 2$

From (i) $(x - \frac{2}{3})$ is a factor

$P(1) = 0$

$\therefore (x-1)$ is a factor

Now $\alpha\beta\gamma = -\frac{2}{6}$

$= -\frac{1}{3}$

$\therefore \frac{2}{3} \times 1 \times \gamma = -\frac{1}{3}$

$\gamma = -\frac{1}{3} \times \frac{3}{2}$

$= -\frac{1}{2}$

$\therefore P(x) = 0$ for $x = -\frac{1}{2}, \frac{2}{3}, 1$ [3]

8.

B	D	B	B	B
---	---	---	---	---

Probability = $\frac{5! \times 4!}{9!}$

$= \frac{1}{126}$

[2]

9. $\int \frac{2}{\sqrt{16-x^2}} dx$

$= 2 \int \frac{dx}{\sqrt{4^2-x^2}}$

$= 2 \sin^{-1} \frac{x}{4} + C$ [2]

(X1) SECTION B

(1) $P(1) = 0 \Rightarrow 1 + 1 + m + n = 0$
 i.e. $m + n = -2$ — (1)

also $P(2) = 11 \Rightarrow 8 + 4 + 2m + n = 11$
 $\therefore 2m + n = -1$ (2)

From (1) + (2)

$$\begin{cases} m = 1 \\ n = -3 \end{cases}$$

$\therefore P(x) = x^3 + x^2 + x - 3$

now $(x-1)$ is a factor.

$\therefore P(x) = (x-1)(x^2 + 2x + 3)$

$$\begin{array}{r} x^2 + 2x + 3 \\ x-1 \overline{) x^3 + 2x^2 + 1x - 3} \\ \underline{x^3 - x^2} \\ 3x^2 + x \\ \underline{3x^2 - 3x} \\ 4x - 3 \end{array}$$

(2.) $f(x) = \sin^{-1}(x-2)$

$\therefore f\left(\frac{3}{2}\right) = \sin^{-1}\left(\frac{3}{2} - 2\right)$
 $= \sin^{-1}\left(-\frac{1}{2}\right)$
 $= \boxed{-\frac{\pi}{6}}$

NB: $|\sin^{-1} \theta| \leq \frac{\pi}{2}$

(3) (i) $\frac{\binom{24}{5}}{\binom{28}{5}} = \frac{253}{585}$

(ii) $\frac{\binom{24}{4} \times \binom{4}{1} + \binom{24}{3} \times \binom{4}{2} + \binom{24}{2} \times \binom{4}{3}}{\binom{28}{5}}$
 $= \frac{2323}{4095}$

(4) $t = \tan x$

$\therefore x = \tan^{-1} t$

$$\left| \frac{dx}{dt} = \frac{1}{1+t^2} \right|$$

(5) (i) $\int \frac{dx}{\sqrt{25-9x^2}} = \int \frac{dx}{\sqrt{9\left(\frac{25}{9} - x^2\right)}}$

$= \frac{1}{3} \int \frac{dx}{\sqrt{\left(\frac{5}{3}\right)^2 - x^2}}$

$= \frac{1}{3} \sin^{-1} \frac{3x}{5} + C$

(ii) $\int \frac{x+1}{x^2+1} dx = \int \frac{x}{x^2+1} dx + \int \frac{dx}{x^2+1}$
 $= \frac{1}{2} \ln(x^2+1) + \tan^{-1} x + C$

$$(6) \int (e^x + e^{-x})^2 dx = \int (e^{2x} + 2e^0 + e^{-2x}) dx$$

$$= \int (e^{2x} + 2 + e^{-2x}) dx$$

$$= \frac{1}{2} e^{2x} + 2x - \frac{1}{2} e^{-2x} + C$$

$$(7) V = \pi \int_0^2 (e^x)^2 dx$$

$$= \pi \int_0^2 e^{2x} dx$$

$$= \pi \left[\frac{1}{2} e^{2x} \right]_0^2$$

$$= \frac{\pi}{2} (e^4 - e^0)$$

$$= \frac{\pi}{2} (e^4 - 1) \text{ m}^3$$

Section C (20 marks)

$$1. \int_0^{\sqrt{2}} \frac{dx}{\sqrt{8-4x}}$$

$$= \int_0^{\sqrt{2}} \frac{dx}{\sqrt{4(2-x)}}$$

$$= \frac{1}{2} \int_0^{\sqrt{2}} \frac{dx}{\sqrt{2-x}}$$

$$= \frac{1}{2} \left[\frac{2\sqrt{2-x}}{\sqrt{2-x}} \right]_0^{\sqrt{2}}$$

$$= \frac{1}{2} \left[\frac{2\sqrt{2-\sqrt{2}}}{\sqrt{2-\sqrt{2}}} - 0 \right]$$

$$= \frac{1}{2} \left[\frac{2\sqrt{2}}{1} \right]$$

$$= \sqrt{2}$$

$$2. y = \ln x \ln x \text{ is curve}$$

$$y' = \ln x \cdot \frac{1}{x} + \frac{1}{x} \ln x$$

$$= \frac{2 \ln x}{x}$$

$$\text{At } P(2, \ln 2) \quad y' = \frac{2 \ln 2}{2} = \ln 2$$

$$\text{is gradient of curve at } P.$$

$$\text{but gradient of base} = \frac{1}{4} \ln 2$$

$$\Rightarrow \frac{1}{4} \ln 2 \neq \ln 2$$

$$\text{Sub into } C \Rightarrow y = \frac{1}{4} \ln 2$$

3. Prove $(9^n - 1)$ is divisible by 8, $n \in \mathbb{Z}^+$

let $n=1$, then $9^1 - 1 = 8$ which is divis. by 8

Assume true for $n=k$

ie. let $9^k - 1 = 8m$ $m \in \mathbb{Z}^+$

let $n=k+1$ Must show $9^{k+1} - 1 = 8p$ $p \in \mathbb{Z}^+$

New $9^{k+1} - 1$

$$= 9^k \cdot 9 - 1$$

$$= 9(9^k - 1) + 8$$

$$= 9(8m) + 8$$

$$= 8(9m + 1)$$

$$= 8p \quad p \in \mathbb{Z}^+$$

True for $n=k+1$

Therefore by the Principle of Mathematical Induction, statement is true for all $n, n \in \mathbb{Z}^+$

3

(2)

At $P(2, \ln 2)$ $y' = \ln 2$ is gradient of curve at $P.$

but gradient of base = $\frac{1}{4} \ln 2$

$$\frac{1}{4} \ln 2 \neq \ln 2$$

\Rightarrow Sub into $C \Rightarrow y = \frac{1}{4} \ln 2$

$y = \frac{1}{4} \ln 2$

3

$\frac{dy}{dx} = 2e^{2x} - 3e^x$ and $y\text{-int} = (0, 3)$
 Then $y = e^{2x} - 3e^x + C$
 $(0, 0) \Rightarrow 0 = 1 - 3 + C$
 $\Rightarrow C = 2$


$y = e^{2x} - 3e^x + 2$
 For x-intercepts, let $y = 0$
 $\Rightarrow e^{2x} - 3e^x + 2 = 0$
 $(e^x - 2)(e^x - 1) = 0$
 $e^x = 1$ or $e^x = 2$
 $x = 0$ or $x = \ln 2$

Area = $\int_0^{\ln 2} (e^{2x} - 3e^x + 2) dx$
 $= \left[\frac{e^{2x}}{2} - 3e^x + 2x \right]_0^{\ln 2}$
 $= \left(\frac{2-6+2\ln 2}{2} - (1-3+0) \right)$
 $= \frac{3}{2} - \ln 4$ square units

5. (i) $y = x \ln x$
 $y' = x^2 \frac{1}{x} + (\ln x) 2x$
 $= x + 2x \ln x$

(ii) Note: $\int (2x + 2x \ln x) dx = x^2 \ln x + C$
 $\Rightarrow \frac{x^2}{2} + 2 \int x \ln x dx = x^2 \ln x + C$
 $\int x \ln x dx = \frac{x^2}{2} \ln x - \frac{x^2}{4} + C$
 $\int (2x \ln x) dx = x^2 \ln x - \frac{x^2}{2} + C$

6. $y = 3 \cos^2 x$
 $\Rightarrow \frac{dy}{dx} = 6 \cos x (-\sin x)$
 $0: -1 \leq 2x \leq 1$ and Range: $0 \leq y \leq 3$
 $0: -\frac{1}{2} \leq x \leq \frac{1}{2}$



(iii) $(e^x - 2)(e^x - 1) = 0$
 $e^x = 1$ or $e^x = 2$
 $x = 0$ or $x = \ln 2$

Area = $\int_0^{\ln 2} (e^{2x} - 3e^x + 2) dx$
 $= \left[\frac{e^{2x}}{2} - 3e^x + 2x \right]_0^{\ln 2}$
 $= \left(\frac{2-6+2\ln 2}{2} - (1-3+0) \right)$
 $= \frac{3}{2} - \ln 4$ square units

SECTION 10

(i) $x^2 = 4ay$
 $\therefore 2x = 4ay$
 $\therefore y' = \frac{x}{2a}$
 At P, $y' = \frac{2ap}{2a} = p$


\therefore Eqn of tangent is
 $y - ap^2 = p(x - 2ap)$
 $y - ap^2 = px - 2ap^2$
 $y = px - ap^2$
 Eqn of A tangent at Q is
 $y = qx - aq^2$

(ii) $px - ap^2 = qx - aq^2$
 $\therefore (p - q)x = ap^2 - aq^2$
 $\therefore (p - q)x = a(p - q)(p + q)$
 $\therefore x = a(p + q)$
 $\therefore y = p(a(p + q)) - ap^2 = apq$
 $\therefore T$ is $(a(p + q), apq)$

(iii) M is $\left(\frac{2ap + 2aq}{2}, \frac{ap^2 + aq^2}{2} \right)$
 $= \left(a(p + q), \frac{a(p^2 + q^2)}{2} \right)$

(iv) R is $\left(\frac{a(p + q) + a(p + q)}{2}, \frac{apq + a(p + q)^2}{2} \right)$
 $= \left(a(p + q), \frac{a(2pq + p^2 + q^2)}{2} \right)$
 $= \left(a(p + q), \frac{a(p + q)^2}{2} \right)$ as given

(v) Let $x = a(p + q)$
 $\therefore y = \frac{x^2}{a} \cdot \frac{1}{4}$
 $x^2 = 4ay$
 $\therefore R$ lies on the parabola $x^2 = 4ay$



10 arrangements.
 \therefore Total arrangements = $10 \times 3 = 30$

$$\begin{aligned} 3 \text{ (i) } (a+b)(a^{n-1} + b^{n-1}) - ab(a^{n-2} + b^{n-2}) \\ = a^n + ab^{n-1} + ba^{n-1} + b^n - a^{n-1}b - ab^{n-1} \\ = a^n + b^n \end{aligned}$$

$$\text{(ii) } x^{k+2} + \frac{1}{x^{k+2}} = \left(x + \frac{1}{x}\right) \left(x^{k+1} + \frac{1}{x^{k+1}}\right) - 1 \cdot \left(x^k + \frac{1}{x^k}\right)$$

$$= 2 \cos \theta \cdot 2 \cos (k+1)\theta - 2 \cos k\theta$$

$$= 2 \cos \theta (2 \cos k\theta \cos \theta - 2 \sin k\theta \sin \theta) - 2 \cos k\theta$$

$$= 4 \cos k\theta \cos^2 \theta - 4 \sin k\theta \sin \theta \cos \theta - 2 \cos k\theta$$

$$= 2 \cos k\theta (2 \cos^2 \theta - 1) - 2 \sin k\theta \sin 2\theta$$

$$= 2 (\cos k\theta \cos 2\theta - \sin k\theta \sin 2\theta)$$

$$= 2 \cos (k+2)\theta \quad \textcircled{2}$$

$$\begin{aligned} \text{(iii) } S(n) &= (x + x^3 + \dots + x^{2n-1}) + \left(\frac{1}{x} + \frac{1}{x^3} + \dots + \frac{1}{x^{2n-1}}\right) \\ &= \frac{\sin 2n\theta}{\sin \theta} \end{aligned}$$

Step 1: show $S(1)$ is true

$$\text{i.e. } x + \frac{1}{x} = \frac{\sin 2\theta}{\sin \theta}$$

$$\text{LHS} = 2 \cos \theta$$

$$\text{RHS} = \frac{2 \sin \theta \cos \theta}{\sin \theta}$$

$$= 2 \cos \theta$$

$\therefore S(1)$ is true

Step 2: Assume $S(k)$ is true

$$\text{i.e. } (x + x^3 + \dots + x^{2k-1}) + \left(\frac{1}{x} + \frac{1}{x^3} + \dots + \frac{1}{x^{2k-1}}\right)$$

$$= \frac{\sin 2k\theta}{\sin \theta}$$

show $S(k+1)$ is true

$$\text{i.e. } (x + x^3 + \dots + x^{2k+1}) + \left(\frac{1}{x} + \frac{1}{x^3} + \dots + \frac{1}{x^{2k+1}}\right)$$

$$= \frac{\sin (2k+2)\theta}{\sin \theta}$$

$$\text{LHS} = \frac{\sin 2k\theta}{\sin \theta} + x^{2k+1} + \frac{1}{x^{2k+1}}$$

$$= \frac{\sin 2k\theta}{\sin \theta} + 2 \cos (2k+1)\theta$$

$$= \frac{\sin 2k\theta + 2 \cos (2k+1)\theta \sin \theta}{\sin \theta}$$

$$= \frac{\sin 2k\theta + 2 \sin \theta (\cos k\theta \cos \theta - \sin k\theta \sin \theta)}{\sin \theta}$$

$$= \frac{\sin 2k\theta + \cos 2\theta \cdot 2 \sin \theta \cos \theta - 2 \sin^2 \theta}{\sin \theta}$$

$$= \frac{\cos 2\theta \cdot \sin 2\theta + \sin 2k\theta (1 - 2 \sin^2 \theta)}{\sin \theta}$$

$$= \frac{\cos 2\theta \sin 2\theta + \sin 2k\theta \cos 2\theta}{\sin \theta}$$

$$= \frac{\sin (2k+2)\theta}{\sin \theta}$$

$$= \text{RHS}$$

\therefore if $S(k)$ is true, $S(k+1)$ is true

Step 3: $S(1)$ is true and $S(k+1)$ is true if $S(k)$ is true.

\therefore By the process of Mathematical

Induction $S(n)$ is true for

all integral $n \geq 1$. $\textcircled{4}$