

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Topic: \_\_\_\_\_

## APPLICATIONS OF INTEGRATION

### Question 1 [3 + 1 + 3 = 7 marks]

The acceleration of a particle undergoing rectilinear motion is given by

$$a = \frac{2}{\sqrt{t+4}} \text{ ms}^{-2}.$$

The particle has a velocity of  $12 \text{ ms}^{-1}$  when  $t = 5$

Find:

- (a) the velocity when  $t = 12$ .

---

---

---

---

---

---

---

---

- (b) if and when the particle is at rest.

---

---

---

---

---

---

---

---

- (c) the distance covered by the particle in the first 5 seconds.

---

---

---

---

---


---

---

---

**Question 2** [1 + 1 + 1 + 2 = 5 marks]

The velocity - time graph below shows the journey taken by two different cyclists, A and B, along the same straight stretch of road.



Use the v - t graph and the fact that they meet after 20 minutes to find:

- (a) the acceleration of A between  $t = 0$  and  $t = 5$ .

---



---



---



---



---

- (b) the displacement of B from his starting position after 20 mins.

---



---



---



---



---

- (c) the total distance travelled by A.

---



---



---



**Question 4** [3 + 3 + 1 + 2 + 1 = 10 marks]

A train slows down with an acceleration which is proportional to its velocity.

- (a) Show that the velocity at any time  $t$  is given by  $v(t) = v_0 e^{kt}$ , where  $v_0$  is the initial velocity.

---

---

---

---

Given that initially the particle has a velocity of  $60 \text{ km h}^{-1}$ , and that the velocity after 5 seconds is  $40 \text{ km h}^{-1}$ , then:

- (b) show that the value of  $k$  is  $\frac{1}{5} \ln \frac{2}{3}$ .

---

---

---

---

Hence, find:

- (c) the velocity after 10 seconds.

---

---

---

- (d) the time taken to reduce the velocity to  $20 \text{ km h}^{-1}$

---

---

---

- (e) the acceleration after 5 seconds.

---

---

---

**Question 5** [2 + 2 = 4 marks]

A particle is moving in a straight line, its velocity at any time  $t$ , is given by

$$v = 6\cos 3t$$

The particle is initially at the origin.

- (a) Find the displacement at any time  $t$ .

- (b) Show that this particle is undergoing Simple Harmonic Motion.

( 7 + 5 + 4 + 10 + 4 = 30 marks )

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Topic: \_\_\_\_\_

**APPLICATIONS OF INTEGRATION****Question 1**

$$(a) \quad v(t) = \int \frac{2dt}{\sqrt{t+4}} = 4\sqrt{t+4} + c \quad [1]$$

$$\text{when } t = 5, v = 12 \Rightarrow c = 0 \quad [1]$$

$$v(12) = 16 \text{ ms}^{-1} \quad [1]$$

$$(b) \quad \text{particle is never at rest since } 4\sqrt{t+4} \neq 0 \quad [1]$$

$$(c) \quad \text{distance travelled} = \int_0^5 4\sqrt{t+4} dt \quad [1]$$

$$= \frac{8}{3}(t+4)^{\frac{3}{2}} \Big|_0^5 \quad [1]$$


$$= 50\frac{2}{3} \text{ m} \quad [1]$$

**Question 2**

$$(a) \quad a = \frac{\text{rise}}{\text{run}} = 0.4 \text{ km min}^{-2} \quad [1]$$

$$(b) \quad x = -10 + 15 = 5 \text{ km to the right of his starting point.} \quad [1]$$

$$(c) \quad \text{dist.} = \frac{5 + 14 + 8}{27} = 27 \text{ km} \quad [1]$$

(d)   $A \quad \quad \quad B \quad \quad \quad 5$   $A \text{ and B finish}$   $[1]$   
initially 22 km apart  $[1]$

**Question 3**

$$V_y = \pi \int_0^{\ln 2} 1 dy - \pi \int_0^{\ln 2} (e^y - 1)^2 dy \quad [1]$$

$$= \pi \int_0^{\ln 2} 1 dy - \pi \int_0^{\ln 2} (e^{2y} - 2e^y + 1) dy$$

$$= \pi \int_0^{\ln 2} (2e^y - e^{2y}) dy \quad [1]$$

$$= \pi \left[ 2e^y - \frac{1}{2}e^{2y} \right]_0^{\ln 2} \quad [1]$$

$$= \pi \left( (4 - 2) - \left( 2 - \frac{1}{2} \right) \right) = \frac{\pi}{2} \quad [1]$$

## Question 4

$$(a) \quad \frac{dv}{dt} = kv \quad [1]$$

$$\ln |v| = kt + c \quad [1]$$

$$v = e^{kt+c} \quad [1]$$

$$v = V_0 e^{kt}$$

$$(b) \quad 40 = 60e^{5k} \quad [1]$$

$$\ln e^{5k} = \ln \frac{2}{3} \quad [1]$$

$$5k = \ln \frac{2}{3} \quad [1]$$

$$k = \frac{1}{5} \ln \frac{2}{3}$$

$$(c) \quad v = 60e^{5 \cdot \frac{1}{5} \ln \frac{2}{3} (10)} = 26.7 \text{ (1dec.pl)} \quad [1]$$

$$(d) \quad 20 = 60e^{5 \cdot \frac{1}{5} \ln \frac{2}{3} t} \quad [1]$$

$$t = 13.6 \text{ (1dec.pl)} \quad [1]$$

$$(e) \quad a = kv(5) = \frac{1}{5} \ln \frac{2}{3} 40 = -3.2 \text{ (1dec.pl)} \quad [1]$$

## Question 5

$$(a) \quad x = 2\sin 3t + c \quad [1]$$

$$\text{when } t = 0, x = 0 \Rightarrow c = 0 \quad [1]$$

$$\therefore x = 2\sin 3t$$

$$(b) \quad a = -18\sin 3t \quad [1]$$

$$= -9 \{2\sin 3t\}$$

$$= -n^2 x \quad [1]$$

$$\therefore \text{S.H.M.}$$

( 7 + 5 + 4 + 10 + 4 = 30 marks )