

--	--	--	--	--	--	--	--

CANDIDATE NUMBER

SYDNEY GRAMMAR SCHOOL



2016 Trial Examination

# FORM VI MATHEMATICS EXTENSION 1

Friday 12th August 2016

**General Instructions**

- Writing time — 2 hours
- Write using black pen.
- Board-approved calculators and templates may be used.

**Total — 70 Marks**

- All questions may be attempted.

**Section I — 10 Marks**

- Questions 1–10 are of equal value.
- Record your answers to the multiple choice on the sheet provided.

**Section II — 60 Marks**

- Questions 11–14 are of equal value.
- All necessary working should be shown.
- Start each question in a new booklet.

**Checklist**

- SGS booklets — 4 per boy
- Multiple choice answer sheet
- Candidature — 109 boys

**Collection**

- Write your candidate number on each answer booklet and on your multiple choice answer sheet.
- Hand in the booklets in a single well-ordered pile.
- Hand in a booklet for each question in Section II, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Write your candidate number on this question paper and hand it in with your answers.
- Place everything inside the answer booklet for Question Eleven.

Examiner  
SO

**SECTION I - Multiple Choice**

Answers for this section should be recorded on the separate answer sheet handed out with this examination paper.

**QUESTION ONE**

What are the asymptotes of  $y = \frac{2x}{(x+3)(x-1)}$ ?

- (A)  $y = 0, x = 1, x = -3$
- (B)  $y = 0, x = -1, x = 3$
- (C)  $y = 2, x = 1, x = -3$
- (D)  $y = 2, x = -1, x = 3$

**QUESTION TWO**

Determine  $\lim_{x \rightarrow 0} \left( \frac{\sin x}{3 \tan x} \right)$ .

- (A) 0
- (B)  $\frac{1}{3}$
- (C) 1
- (D) 3

**QUESTION THREE**

What is the domain of  $f(x) = e^{-\frac{1}{x}}$ ?

- (A)  $x > 0$
- (B)  $x \geq 0$
- (C)  $x \neq 0$
- (D) all real  $x$

**QUESTION FOUR**

What is the value of  $\sin(\tan^{-1} a)$ ?

- (A)  $\frac{a}{\sqrt{1-a^2}}$
- (B)  $\frac{1}{\sqrt{1-a^2}}$
- (C)  $\frac{1}{\sqrt{1+a^2}}$
- (D)  $\frac{a}{\sqrt{1+a^2}}$

**QUESTION FIVE**

The monic quadratic equation with roots  $m + n$  and  $m - n$  is:

- (A)  $x^2 - 2mx + m^2 - n^2 = 0$
- (B)  $x^2 + 2mx + m^2 - n^2 = 0$
- (C)  $x^2 - 2mx + n^2 - m^2 = 0$
- (D)  $x^2 + 2mx + n^2 - m^2 = 0$

**QUESTION SIX**

A function is defined by the following rule:

$$f(x) = \begin{cases} \sin^{-1} x, & \text{for } -1 \leq x < 0 \\ \cos^{-1} x & \text{for } 0 \leq x \leq 1 \end{cases}$$

What is the value of  $f(-\frac{1}{2}) + f(0)$ ?

- (A)  $-\frac{\pi}{6}$
- (B)  $\frac{\pi}{6}$
- (C)  $\frac{\pi}{3}$
- (D)  $\frac{2\pi}{3}$

**QUESTION SEVEN**


The range  $R$  of any particle projected from a point on a level plane at an angle of  $\alpha$  to the horizontal with initial speed  $v$  is given by  $R = \frac{v^2 \sin 2\alpha}{g}$ .

A particle is projected at  $50^\circ$  to the horizontal. What other angle of projection would give the same range for this particle?

- (A)  $25^\circ$
- (B)  $40^\circ$
- (C)  $80^\circ$
- (D)  $100^\circ$

Examination continues overleaf ...

**QUESTION EIGHT**



The points  $A, B$  and  $C$  lie on a circle with centre  $O$ . If  $\angle ABC = 130^\circ$ , what is the size of  $\angle AOC$ ?

- (A)  $50^\circ$
- (B)  $65^\circ$
- (C)  $100^\circ$
- (D)  $260^\circ$

**QUESTION NINE**

A particle is moving in simple harmonic motion with period 4 and amplitude 3. Which of the following is a possible equation for the velocity of the particle?

- (A)  $v = \frac{3\pi}{2} \cos \frac{\pi t}{2}$
- (B)  $v = 3 \cos \frac{\pi t}{2}$
- (C)  $v = \frac{3\pi}{4} \cos \frac{\pi t}{4}$
- (D)  $v = 3 \cos \frac{\pi t}{4}$

**QUESTION TEN**

Which of the following is a necessary condition if  $a^2 > b^2$ ?

- (A)  $a > b$
- (B)  $a < b < 0$
- (C)  $a > 0 > b$
- (D)  $|a| > |b|$

\_\_\_\_\_ End of Section I \_\_\_\_\_

Examination continues next page ...

SECTION II - Written Response

Answers for this section should be recorded in the booklets provided.

Show all necessary working.


Start a new booklet for each question.

QUESTION ELEVEN (15 marks) Use a separate writing booklet. Marks

- (a) Find the value of  $\sin^{-1}(\sin \frac{3\pi}{5})$ . 1
- (b) Let  $A = (4, -3)$  and  $B = (8, 5)$ . The interval  $AB$  is divided internally in the ratio  $3 : 1$  by the point  $P(x, y)$ . Find the values of  $x$  and  $y$ . 2
- (c) Solve  $\frac{5}{3x-2} > 2$ . 3
- (d) The acute angle between the two lines  $y = \frac{1}{2}x + 1$  and  $y = mx + 3$  is  $\frac{\pi}{4}$ . Find all possible values of the constant  $m$ . 3
- (e) Find the general solution of  $\cos 2x - \cos x = 2$ . 3
- (f) Find the term independent of  $x$  in the expansion of  $(x^2 - \frac{2}{x})^{12}$ . 3

QUESTION TWELVE (15 marks) Use a separate writing booklet. Marks

- (a) The polynomial  $P(x) = 2x^3 + x^2 + ax + 6$  has a zero at  $x = 2$ .
  - (i) Determine the value of  $a$ . 1
  - (ii) Find the linear factors of  $P(x)$ . 2
  - (iii) Hence, or otherwise, solve  $P(x) \geq 0$ . 1
- (b) Integrate  $\int_0^{\frac{1}{\sqrt{3}}} \frac{\sin(\tan^{-1} x)}{1+x^2} dx$  using the substitution  $u = \tan^{-1} x$ . 3
- (c) 2




The diagram above shows a hot air balloon at point  $H$  with altitude  $800\text{m}$ . The passengers in the balloon can see a barn and a dam below, at points  $B$  and  $D$  respectively. Point  $C$  is directly below the hot air balloon. From the hot air balloon's position, the barn has a bearing of  $250^\circ$  and the dam has a bearing of  $130^\circ$ , and  $\angle BCD = 120^\circ$ . The angles of depression to the barn and the dam are  $50^\circ$  and  $30^\circ$  respectively.

How far is the barn from the dam, to the nearest metre?

(d) Prove by induction that  $(x + y)$  is a factor of  $x^{2n} - y^{2n}$ , for all integers  $n \geq 1$ . 3


(e) 3



The diagram above shows a vessel in the shape of a cone of radius 3 cm and height 4 cm. Water is poured into it at the rate of  $10 \text{ cm}^3/\text{s}$ . Find the rate at which the water level is rising when the depth is 2 cm.

QUESTION THIRTEEN (15 marks) Use a separate writing booklet. Marks

(a)



Two circles  $C_1$  and  $C_2$  intersect at  $A$  and  $B$ . A line through  $A$  meets the circles at  $P$  and  $Q$  respectively. A tangent is drawn from an external point  $T$  to touch the circle  $C_1$  at  $P$ . The line  $TQ$  intersects  $C_2$  at  $R$ .

(i) Given  $\angle XPB = \alpha$ , show that  $\angle BRQ = 180^\circ - \alpha$ , giving reasons. 2

(ii) Hence show that  $PTRB$  is a cyclic quadrilateral. 1

(b) Consider the parabola  $x^2 = 4ay$  with focus  $S$ . The normal at  $P(2ap, ap^2)$  meets the  $y$ -axis at  $R$  and  $\triangle SPR$  is equilateral.

(i) Show that the equation of the normal at  $P$  is  $x + py = 2ap + ap^3$ . 1

(ii) Write down the coordinates of  $R$ . 1

(iii) Prove that  $SP$  is equal in length to the latus rectum, that is  $4a$  units. 3


(c) (i) Show that  $\frac{d}{dx}(x \ln x) = 1 + \ln x$ . 1

(ii) A particle is moving in a straight line. At time  $t$  seconds its position is  $x$  cm and its velocity is  $v$  cm/s. Initially  $x = 1$  and  $v = 2$ . The acceleration  $a$  of the particle is given by the equation

$$a = 1 + \ln x.$$

Find the velocity  $v$  in terms of  $x$ . Be careful to explain why  $v$  is always positive. 2

(d)



The circle above has radius 1 unit and the major arc joining  $A$  and  $B$  is twice as long as the chord  $AB$ . The point  $M$  lies on  $AB$  such that  $AB \perp OM$ . Let  $\angle AOM = \theta$  where  $0 < \theta < \frac{\pi}{2}$ .

- (i) Show that the length of the major arc satisfies the equation 1  

$$\pi - \theta = 2 \sin \theta.$$
- (ii) Let  $\theta_0 \doteq 1.5$  be a first approximation of  $\theta$ . Use two applications of Newton's method to find a better approximation of  $\theta$ . 2
- (iii) Use your answer to part (ii) to find the approximate length of the chord  $AB$ . 1

Examination continues overleaf ...

QUESTION FOURTEEN (15 marks) Use a separate writing booklet. Marks

- (a) The mass  $M$  of a radioactive isotope is given by the equation  $M = M_0 e^{-kt}$ , where  $M_0$  is the initial mass and  $k$  is a constant. The mass satisfies the equation  $\frac{dM}{dt} = -kM$ .
  - (i) If the half-life of this radioactive isotope is  $T$ , show that  $k = \frac{\log_e 2}{T}$ . 1
  - (ii) A naturally occurring rock contains two radioactive isotopes  $X$  and  $Y$ . The half-lives of isotope  $X$  and isotope  $Y$  are  $T_X$  and  $T_Y$  respectively, where  $T_X > T_Y$ . Initially the mass of isotope  $Y$  is four times that of isotope  $X$ . 3

Show that the rock will contain the same mass of both isotopes at time

$$\frac{2T_X T_Y}{T_X - T_Y}.$$

- (b) Sketch the graph of  $y = \frac{|x|}{x}$ . 1
- (c) Consider the function  $f(x) = \sin^{-1} \left( \frac{x^2 - 1}{x^2 + 1} \right)$ .
  - (i) Find the domain of  $f(x)$ . 1
  - (ii) Show that  $f'(x) = \frac{2x}{|x|(x^2 + 1)}$ . 2
  - (iii) Determine the values of  $x$  for which  $f(x)$  is increasing. 1
  - (iv) Using part (b), explain the behaviour of  $f'(x)$  as  $x \rightarrow 0^+$  and  $x \rightarrow 0^-$ . 1
  - (v) Draw a neat sketch of  $y = f(x)$ , indicating any intercepts with the axes and any asymptotes. 2
  - (vi) Give the largest possible domain containing  $x = 1$  for which  $f(x)$  has an inverse function. Let this inverse function be  $f^{-1}(x)$ . 1
  - (vii) Sketch  $y = f^{-1}(x)$  on your original graph. 1
  - (viii) Find the equation of  $f^{-1}(x)$ . 1

————— End of Section II —————

END OF EXAMINATION

# EXTENSION MATHEMATICS SOLUTIONS


## MULTIPLE CHOICE

1. Vertical asymptotes:  $(x+3)(x-1) = 0$   
 $x = -3$  and  $x = 1$       A

Horizontal asymptote:  $y = 0$

2.  $\lim_{x \rightarrow 0} \frac{\sin x}{x} \times \frac{x}{\tan x} \times \frac{1}{3} = \frac{1}{3}$       B

3.  $f(x) = e^{-\frac{1}{x}}$      $x \neq 0$       C



5.  $(x - (m+n))(x - (m-n)) = 0$   
 $x^2 - 2mx + m^2 - n^2 = 0$       A

6.  $f(-\frac{1}{2}) + f(0)$   
 $= \sin^{-1}(-\frac{1}{2}) + \cos^{-1}(0)$   
 $= -\frac{\pi}{6} + \frac{\pi}{2}$       C  
 $= \frac{\pi}{3}$

7.  $\sin(2 \times 50^\circ) = \sin 100^\circ = \sin 80^\circ = \sin(2 \times 40^\circ)$       B

8. Reflex  $\angle AOC = 260^\circ$  (angle at the centre is twice the angle at the circumference)  
 $\therefore \angle AOC = 100^\circ$       C

9.  $T = \frac{2\pi}{\omega}$        $x = 3 \sin \frac{\pi t}{2}$   
 $= \frac{\pi}{2}$        $V = \frac{3\pi}{2} \cos \frac{\pi t}{2}$       A

10. A sufficient  
 B neither necessary nor sufficient  
 C sufficient  
 D necessary and sufficient

1	2	3	4	5	6	7	8	9	10
A	B	C	D	A	E	B	C	A	D

Q11

(a)  $\sin^{-1}(\sin(\frac{3\pi}{5}))$   
 $= \sin^{-1}(\sin(\frac{2\pi}{5}))$   
 $= \frac{2\pi}{5}$  ✓

(b) A B  
 $(4, -3)$   $(8, 5)$   
 3 ; 1

$x = \frac{3 \times 8 + 1 \times 4}{4} = \frac{28}{4} = 7$  ✓  
 $y = \frac{3 \times 5 + 1 \times 3}{4} = \frac{18}{4} = 3$  ✓

So  $(x, y) = (7, 3)$

(c)  $\frac{5}{3x-2} > 2$


$5(3x-2) > 2(3x-2)^2$  ✓  $x + \frac{2}{3}$

$2(3x-2)^2 - 5(3x-2) < 0$

$(3x-2)(6x-9) < 0$  ✓

$3(3x-2)(2x-3) < 0$

So  $\frac{2}{3} < x < \frac{3}{2}$  ✓



(d)  $\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$   
 $1 = \left| \frac{m_1 - \frac{1}{2}}{1 + \frac{m_1}{2}} \right|$  ✓

$1 + \frac{m_1}{2} = m_1 - \frac{1}{2}$  or  $1 + \frac{m_1}{2} = -m_1 + \frac{1}{2}$   
 $-\frac{m_1}{2} = -\frac{3}{2}$  or  $\frac{3m_1}{2} = -\frac{1}{2}$   
 $m_1 = 3$  ✓  $m_1 = -\frac{1}{3}$  ✓

So the possible values of the constant  $m$  are 3 and  $-\frac{1}{3}$

(e)  $\cos 2x - \cos x = 2$

$2\cos^2 x - 1 - \cos x - 2 = 0$  ✓

$2\cos^2 x - \cos x - 3 = 0$

$(2\cos x - 3)(\cos x + 1) = 0$

$\cos x = \frac{3}{2}$

$\cos x = -1$  ✓

↓  
no solution

$x = -\pi + 2n\pi$ , where  $n$  is an integer ✓

(f) Let the term be  ${}^{12}C_k (x^2)^{k-k} \left(-\frac{2}{x}\right)^k$

$= {}^{12}C_k \cdot x \cdot x^{24-2k} \times (-1)^k \times 2^k \times x^{-k}$

$= {}^{12}C_k \times x^{24-3k} \times (-1)^k \times 2^k$  ✓

For the term independent of  $x$ ,  $24 - 3k = 0$   
 $k = 8$  ✓

So  ${}^{12}C_8 \times x^0 \times (-1)^8 \times 2^8$

$= {}^{12}C_8 \times 2^8$  ✓

$= 495 \times 256$

$= 126720$

QN 12

(a)(i)  $P(x) = 0$

$0 = 16x + 4x + 2a + 6$

$2a = -20$

$a = -10$


(ii)  $\frac{2x^2 + 5x - 3}{x-2} \cdot \frac{2x^2 + x^2 - 13x + 6}{2x^3 - 4x^2}$

$2x^2 + 5x - 3 = (2x-1)(x+3)$

so  $P(x) = (x-2)(2x-1)(x+3)$

$\frac{5x^2 - 13x}{5x^2 - 10x}$   
 $\frac{-3x + 6}{-3x + 6}$

(iii)  $P(x) \geq 0$   
 $-3 \leq x \leq \frac{1}{2}$  or  $x \geq 2$



(b)  $\int_0^{\frac{1}{\sqrt{3}}} \frac{\sin(\tan^{-1}x)}{1+x^2} dx$

$u = \tan^{-1}x$   
 $du = \frac{1}{1+x^2} dx$   
 when  $x = \frac{1}{\sqrt{3}}$   $u = \frac{\pi}{6}$   
 $x = 0$   $u = 0$


$= \int_0^{\frac{\pi}{6}} \sin u du$

$= -[\cos u]_0^{\frac{\pi}{6}}$

$= -\left(\frac{\sqrt{3}}{2} - 1\right)$

$= \frac{1 - \frac{\sqrt{3}}{2}}{1}$

(c)



In  $\triangle BCH$ ,  $\tan 50^\circ = \frac{800}{BC}$

$BC = 800 \cot 50^\circ$

Similarly in  $\triangle DCH$ ,  $\tan 30^\circ = \frac{800}{DC}$

$DC = 800 \cot 30^\circ$

By the cosine rule:

$BD^2 = (800 \cot 50^\circ)^2 + (800 \cot 30^\circ)^2 - 2 \cdot 800 \cot 50^\circ \cdot 800 \cot 30^\circ \cos 120^\circ$

$= 800^2 \times 5.157$

$= 3300488$

$= 1816.722...$

$= 1817m$

(d) Prove  $x^{2n} - y^{2n}$  has  $(x+y)$  as a factor for all integers  $n > 1$ .

① For  $n=1$ :  $x^2 - y^2 = (x+y)(x-y)$ , which has  $(x+y)$  as a factor.

② Assume true for  $n=k$ :

$x^{2k} - y^{2k} = m(x+y)$  where  $m$  is an expression in  $x$  &  $y$

③ Prove true for  $n=k+1$ :

From ②  $x^{2k+2} - y^{2k+2} = m(x+y) + y^{2k}$

$x^{2k+2} - y^{2k+2}$

$= x \cdot x^{2k} - y \cdot y^{2k}$

$= x^2 [m(x+y) + y^{2k}] - y^2 \cdot y^{2k}$


$= mx^2(x+y) + x^2 y^{2k} - y^2 \cdot y^{2k}$

$= mx^2(x+y) + y^{2k}(x^2 - y^2)$

$= x+y [mx^2 + y^{2k}(x-y)]$



So from parts (2) & (3) and by mathematical induction  
 $(x+y)$  is a factor of  $x^{2n} - y^{2n}$  for all integers  $n \geq 1$ .



$$\frac{r}{h} = \frac{3}{4}$$

$$r = \frac{3h}{4}$$

$$V = \frac{1}{3} \pi \left( \frac{3h}{4} \right)^2 h$$

$$= \frac{3h^3 \pi}{16}$$

$$\frac{dV}{dh} = \frac{9h^2 \pi}{16}$$

$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}$$

$$10 = \frac{9 \times 4 \times \pi}{16} \times \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{40}{9\pi} \text{ cm/s}$$

So the water is rising at a rate of  $\frac{40}{9\pi}$  cm/s when the depth of the cone is 2cm.

Q13

(a)(i)  $\angle XPB = \alpha$ . Join points A and B, and B and R.

$\angle PAB = \alpha$  (alternate segment theorem) ✓

$\angle BAO = 180^\circ - \alpha$  (straight line)

$\angle BRQ = 180^\circ - \alpha$  (angles standing on the same arc) ✓

(ii)  $\angle TRB = \alpha$  (straight angle)

$\angle XPT = \angle TRB$

$\therefore$  PTRB is a cyclic quadrilateral

(Exterior angle is equal to the opposite interior angle) ✓

(b)(i)  $\frac{dy}{dx} = \frac{2ap}{2a}$

$= p$

so gradient of the normal is  $-\frac{1}{p}$

equation of the normal:


$$y - ap^2 = -\frac{1}{p}(x - 2ap)$$

$$py - ap^3 = -x + 2ap$$

$$x + py = 2ap + ap^3$$

(ii)  $R(0, 2a + ap^2)$

(iii) Let F be foot of the perpendicular from P to the y-axis.



Then  $FR = FS = 2a$  ✓

( $\Delta SRP$  is isosceles with altitude  $FP$ )

Hence  $SR = 4a$  and since  $\Delta SPR$  is equilateral

$$SP = SR = 4a$$

= length of latus rectum. ✓✓

(c) (i)  $\frac{d}{dx}(x \ln x) = \ln x + \frac{1}{x} \times x$   
 $= \ln x + 1$  ✓

(ii)  $a = 1 + \ln x$

$$\frac{1}{2} v^2 = x \ln x + c$$

at  $v=2, x=1$  so  $c=2$

$$\frac{1}{2} v^2 = x \ln x + 2$$

$$v^2 = 2x \ln x + 4$$

$$v = \pm \sqrt{2x \ln x + 4}$$

$\ln x$  is always increasing

so at  $t=0, a > 0$

so  $v$  is always positive ✓

$$v = \sqrt{2x \ln x + 4}$$


(d) (i) Let  $AM = x$

chord  $AB = 2x$   
 major arc  $AB = 4x$

$x = \sin\theta$   
 so  $AB = 2\sin\theta$

Major arc  $AB = 2\pi - 2\theta$

so  $2\pi - 2\theta = 4\sin\theta$   
 $\pi - \theta = 2\sin\theta$  ✓



(ii)  $2\sin\theta + \theta - \pi = 0$

So  $f(\theta) = 2\sin\theta + \theta - \pi$

$f'(\theta) = 2\cos\theta + 1$  ✓

$\theta_1 = \theta_0 - \frac{f(\theta)}{f'(\theta)}$

$= 1.5 - \frac{(2\sin\theta + \theta - \pi)}{2\cos\theta + 1}$

$= 1.19$

$\theta_2 = 1.24433...$  ✓

(iii)  $AB = 2\sin\theta$

$= 2 \times \sin(1.24433...)$

$= 1.89436...$  ✓

**Q14**

(a) (i)  $\frac{1}{2} M_0 = M_0 e^{-kT}$

$\frac{1}{2} = e^{-kT}$

$\ln\left(\frac{1}{2}\right) = -kT$

$k = \frac{\ln 2}{T}$  ✓

(ii)  $X = X_0 e^{-k_x t}$

$Y = Y_0 e^{-k_y t} = 4X_0 e^{-k_y t}$

$4X_0 e^{-k_y t} = X_0 e^{-k_x t}$  ✓

$\frac{4}{e^{k_y t}} = \frac{1}{e^{k_x t}}$

$4 = \frac{e^{k_x t}}{e^{k_y t}}$

$4 = e^{t(k_x - k_y)}$


$2\ln 2 = t(k_x - k_y)$

$2\ln 2 = t\left(\frac{\ln 2}{T_y} - \frac{\ln 2}{T_x}\right)$  ✓ (using part (i))

$2 = t\left(\frac{T_x - T_y}{T_x T_y}\right)$

$t = \frac{2T_x T_y}{T_x - T_y}$  ✓

(b)



(c) (i)  $f(x) = \sin^{-1}\left(\frac{x^2-1}{x^2+1}\right)$

Domain:  $-1 \leq \left(\frac{x^2-1}{x^2+1}\right) \leq 1$

$$-x^2-1 \leq x^2-1 \leq x^2+1$$

$$\begin{aligned} -x^2-1 &\leq x^2-1 & \text{and} & & x^2-1 &\leq x^2+1 \\ -2x^2 &\leq 0 & & & 0 &\leq 2 \\ -x^2 &\leq 0 & & & \text{true for all } x & \end{aligned}$$

So domain is all real  $x$  ✓

(ii) Let  $y = \sin^{-1}u$

$$\frac{dy}{du} = \frac{1}{\sqrt{1-u^2}}$$

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$= \frac{1}{\sqrt{1-\left(\frac{x^2-1}{x^2+1}\right)^2}} \times \frac{4x}{(x^2+1)^2}$$

$$= \frac{4x}{(x^2+1) \times 4x}$$

$$= \frac{4x}{\sqrt{x^4+2x^2+1-(x^4-2x^2+1)}} \times (x^2+1)^2$$

$$= \frac{4x}{\sqrt{4x^2} \times (x^2+1)}$$

$$\begin{aligned} u &= x^2-1 & v &= x^2+1 \\ u' &= 2x & v' &= 2x \end{aligned}$$

$$\begin{aligned} \frac{du}{dx} &= \frac{2x(x^2+1) - 2x(x^2-1)}{(x^2+1)^2} \\ &= \frac{2x(x^2+1-x^2+1)}{(x^2+1)^2} \\ &= \frac{4x}{(x^2+1)^2} \end{aligned}$$

$$= \frac{4x}{\sqrt{4x^2} \times (x^2+1)}$$

$$= \frac{2|x| \times (x^2+1)}{|x| \times (x^2+1)} \quad (\text{by definition } \sqrt{x^2} = |x|)$$


$$= \frac{2x}{|x| \times (x^2+1)} \quad \checkmark$$

(ii) Increasing when  $\frac{dy}{dx} > 0$   
 $x > 0$  ✓

(iv)  $\lim_{x \rightarrow 0^+} f'(x) = 2$

$$\lim_{x \rightarrow 0^-} f'(x) = -2 \quad \checkmark$$

The gradient approaches +2 from below and -2 from above



$x$ -intercept at  $f(x) = 0$

$$\begin{aligned} x^2-1 &= 0 \\ x &= \pm 1 \end{aligned}$$

Asymptotes

$$\text{as } x \rightarrow \infty \quad f(x) \rightarrow \sin^{-1}1 \rightarrow \frac{\pi}{2}$$

$$\text{as } x \rightarrow -\infty \quad f(x) \rightarrow \sin^{-1}(-1) \rightarrow -\frac{\pi}{2}$$

$$\text{Horizontal asymptote: } y = \frac{\pi}{2}$$

$$\text{Horizontal asymptote: } y = -\frac{\pi}{2}$$

$$(vi) \quad -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \quad \checkmark$$

(vii) See graph  $\checkmark$

$$(viii) \quad x = \sin^{-1}\left(\frac{y^2-1}{y^2+1}\right)$$

$$\sin x = \frac{y^2-1}{y^2+1}$$

$$\sin x (y^2+1) = y^2-1$$

$$y^2 \sin x + \sin x = y^2-1$$

$$y^2 (1-\sin x) = 1+\sin x$$

$$y^2 = \frac{1+\sin x}{1-\sin x}$$

$$y = \sqrt{\frac{1+\sin x}{1-\sin x}} \quad \checkmark$$

$$\therefore f^{-1}(x) = \sqrt{\frac{1+\sin x}{1-\sin x}}$$