

Name: Maths Class:

SYDNEY TECHNICAL HIGH SCHOOL



Year 12 Mathematics Extension 1

HSC Course
Assessment 2
March, 2016

Time allowed: 90 minutes

General Instructions:

- Marks for each question are indicated on the question.
- Approved calculators may be used
- All necessary working should be shown
- Full marks may not be awarded for careless work or illegible writing
- *Begin each question on a new page*
- Write using black or blue pen
- All answers are to be in the writing booklet provided
- A BOSTES Reference Sheet is provided at the rear of this Question Booklet, and may be removed.


Section I Multiple Choice
Questions 1-6
6 Marks

Section II Questions 7-12
60 Marks

QUESTION 1

O is the centre. Diagram is not to scale.

What is the size of x in degrees?




- A. 110 B. 120 C. 130 D. 140

QUESTION 2

O is the centre. Diagram is not to scale.


What is the size of x in degrees?



- A. 30 B. 45 C. 60 D. 75

QUESTION 3

Diagram is not to scale. What is the length of x in units?




- A. 6 B. 6.25 C. 6.5 D. 6.75

QUESTION 4

Which of the following is not a possible

function for this curve?

- A. $y = \cos 2x$
 B. $y = \sin 2\left(x - \frac{3\pi}{4}\right)$
 C. $y = \sin\left(2x + \frac{\pi}{4}\right)$
 D. $y = -\cos 2\left(x - \frac{\pi}{2}\right)$



QUESTION 5


For which values of x is the curve $y = x^3 - 3x^2$ both concave up and decreasing?

- A. $x > 1$ B. $1 < x < 2$ C. $0 < x < 2$ D. $0 < x < 1$

QUESTION 6

The graph of a derivative function is shown.

At which x value on the original curve $y = f(x)$ will a horizontal point of inflexion occur?



- A. $x = A$
 B. $x = B$
 C. $x = C$
 D. $x = D$

QUESTION 7 (10 marks) Start a new page.

- a) Write the exact value of $\sec \frac{5\pi}{4}$ 1
 b) i) Differentiate $x(2x + 1)^3$ 1
 ii) Find the equation of the tangent to the curve $y = x(2x + 1)^3$ at the point where $x = -1$. 2
 c) Differentiate $\tan(\sin 2x)$ 2
 d) Solve $3\cos^2 x = \cos x$ for $0 \leq x \leq 2\pi$. Give solutions correct to 2 decimal places where necessary. 3
 e) Find $\int \frac{x^2+1}{x^2} dx$ 1


QUESTION 8 (10 marks) Start a new page.

- a) Find $\int (mx - k)^3 dx$ 1
 b) The curve $y = ax^2 + bx$ has a gradient of 6 at (2,0). Find the values of a and b . 2

c) The area of the shaded segment is 100 cm^2 and

subtends an angle of $\frac{5\pi}{6}$ radians at the centre.

Find the radius, correct to 1 decimal place.



d) Given the curve $y = 1 - 2 \cos \frac{\pi x}{5}$, x in radians.

- i) Find the period of this curve. 1
 ii) Find x intercepts for one period, $x \geq 0$. 2
 iii) Sketch the curve for one period, $x \geq 0$. Show x and y intercepts. 2

QUESTION 9 (10 marks) Start a new page.

a) Using a neat diagram and a ruler, find an approximate solution to the equation

$$\sin \frac{x}{2} = 1 - \frac{x}{3}$$

b) For the curve $y = \frac{-3x}{(x-1)^2}$:


- i) What is the equation of the vertical asymptote? 1
 ii) Find the stationary point and determine its nature. 3
 iii) Examine the behaviour of the curve as $x \rightarrow \pm\infty$ 1
 iv) Neatly sketch the curve. Use a ruler and show relevant features. 2
 v) Indicate, with an arrow, the location of a point of inflexion on the curve. 1

Do not find its coordinates.


QUESTION 10 (11 marks) Start a new page.

a) Evaluate $\int_{-\sqrt{7}}^0 x\sqrt{16-x^2} dx$, using the substitution $u = 16 - x^2$ 3

b) Find the shaded area:



- c) Consider the largest cylinder with height h units and radius r units that can fit inside a sphere of fixed radius a units.




- i) Show that the cylinder has volume $V = \frac{\pi}{4}(4a^2h - h^3)$. 2
- ii) Prove that the maximum volume of the cylinder is $\frac{4\pi a^3}{3\sqrt{3}}$. 3

QUESTION 11 (10 marks) Start a new page.

- a) Use Mathematical Induction to prove that $9^n - 5^n$ is always divisible by 4, for all positive integers n . 3
- b) i) On the same axes, sketch the curves $y = x^2$ and $x = y^2$. Show points of intersection. 1
- ii) Find the magnitude of the area bounded by the two curves. 3
- c) The area between $y = 1 - x^2$ and the coordinate axes is rotated about the y axis. 3
- Find the volume of the generated solid.

QUESTION 12 (9 marks) Start a new page.


- a) A, B, C, D are points on the circle, centre O.
- $AB = CD$



- i) Neatly redraw the diagram into your answer booklet.
- ii) If $\angle DAC = x$ degrees, find the size of $\angle BAC$ in terms of x . Give reasons.

3

- b) The curve shows a function $y = f(x)$.




Neatly sketch a possible curve for $y = f''(x)$.

Show corresponding positions P to T on your graph.

3

- c) The shaded area below is rotated about the x axis.



Use Simpson's Rule and five function values to estimate the volume of the generated solid. Give your answer correct to one decimal place.

3

END OF EXAM

SOLUTIONS

- ① D ② B ③ B ④ C ⑤ B ⑥ D

1) a) $\frac{1}{\cos 225^\circ} = \frac{-1}{\cos 45^\circ} = -\sqrt{2}$

b) i) $y' = 1(2x+1)^3 + 3(2x+1)^2 \times 2 \times 2 = (2x+1)^3 + 6x(2x+1)^2$

ii) $x = -1 \Rightarrow y' = -1 - 6(1) = -7$
 $m_T = -7$ and $(-1, 1)$

\therefore tangent is $y - 1 = -7(x + 1)$
 $y = -7x - 6$

c) $y' = \sec^2(\sin 2x) \times \cos 2x \times 2 = 2 \sec^2(\sin 2x) \cos 2x$

d) $3 \cos^2 x - \cos x = 0$
 $\cos x(3 \cos x - 1) = 0$
 $\cos x = 0$ or $\frac{1}{3}$
 $x = \frac{\pi}{2}, \frac{3\pi}{2}, 1.23, 5.05$

e) $\int (1 + x^{-2}) dx = x - \frac{1}{x} + c$

8) a) $(mx - k)^4 + c$
 $4cm$

b) $y' = 2ax + b$. At $x=2, y'=6$
 $\therefore 6 = 4a + b$ — (1)

At $x=2, y=0$
 $\therefore 0 = 4a + 2b$ — (2)

② - ① gives $b = -6$



Sub in ② gives $4a - 12 = 0$
 $a = 3$

c) $\frac{1}{2} r^2 (\theta - \sin \theta) = 100$
 $r^2 (5\frac{\pi}{6} - \frac{1}{2}) = 200$
 $r^2 = 200 \times \frac{6}{5\pi - 3}$

$r \doteq 9.7$ (1 dec.)

d) i) period = $\frac{2\pi}{\frac{\pi}{5}} = 10$

d) ii) $-2 \cos \frac{\pi x}{5} = 0$
 $\cos \frac{\pi x}{5} = \frac{1}{2}$
 $\frac{\pi x}{5} = \frac{\pi}{3}, \frac{5\pi}{3}$
 $x = \frac{5}{3}$ or $\frac{25}{3}$



b) i) $x = 1$ $x = 1.4, 1.5, 1.6$ or $\frac{\pi}{2}$

ii) S.P. when $y' = \frac{-3(x-1)^2 - 2(x-1)(-3x)}{(x-1)^4} = 0$

$\therefore -3(x^2 - 2x + 1) + 6x^2 - 6x = 0$
 $-3x^2 + 6x - 3 + 6x^2 - 6x = 0$
 $3x^2 - 3 = 0$
 $3(x+1)(x-1) = 0$

$x = \pm 1$ (but $x \neq 1$).
 $\therefore x = -1$ only

$y' = \frac{3(x-1)(x+1)}{(x-1)^4 + 3} = \frac{3(x+1)}{(x-1)^3}$


x	-1.1	-1	-0.9
y'	+	0	-

\therefore max T.P. at $(-1, \frac{3}{4})$

$$(iii) y = \frac{-3x}{x^2 - 2x + 1}$$

$$= \frac{-3x}{(x - \frac{1}{2})^2}$$

As $x \rightarrow \pm \infty$,
 $y \rightarrow 0$



(10) a) $\int_{-\sqrt{5}}^0 x \sqrt{16-x^2} dx = \int_9^{16} x \sqrt{u} \frac{du}{-2x}$

$u = 16 - x^2, x = -\sqrt{u}$
 $\frac{du}{dx} = -2x, u = 9$
 $dx = \frac{du}{-2x}, u = 16$

$$= -\frac{1}{2} \int_9^{16} u^{1/2} du$$

$$= -\frac{1}{2} \times \frac{2}{3} \left[u^{3/2} \right]_9^{16}$$

$$= -\frac{1}{3} (64 - 27)$$

$$= -\frac{37}{3}$$

b) Area = square - area to x axis

$$= (1 \times 1) - \int_0^1 (2x - x^2) dx$$

$$= 1 - \left[x^2 - \frac{x^3}{3} \right]_0^1$$

$$= 1 - \left[\left(1 - \frac{1}{3}\right) - (0 - 0) \right]$$

$$= 1 - \frac{2}{3}$$

$$= \frac{1}{3} u^2$$

c) i) $V = \pi r^2 h$
and $a^2 = \left(\frac{h}{2}\right)^2 + r^2$
 $\therefore r^2 = a^2 - \frac{h^2}{4}$

$$\therefore V = \pi \left(\frac{4a^2 - h^2}{4} \right) h$$

$$= \frac{\pi}{4} (4a^2 h - h^3) u^3$$

as reqd.

c) ii) max vol. when $\frac{dV}{dh} = 0$

$$\frac{dV}{dh} = \frac{\pi}{4} (4a^2 - 3h^2) = 0$$

$$3h^2 = 4a^2$$

$$h^2 = \frac{4a^2}{3}$$

$$h = \frac{2a}{\sqrt{3}}$$

Prove max. vol.
 $V'' = \frac{\pi}{4} (-6h) < 0$ for all h .
 \therefore max. vol. occurs when $h = \frac{2a}{\sqrt{3}}$

and $V_{max} = \frac{\pi}{4} \left(4a^2 \times \frac{2a}{\sqrt{3}} - \frac{8a^3}{3\sqrt{3}} \right)$

$$= \frac{\pi}{4} \left(\frac{8a^3}{\sqrt{3}} - \frac{8a^3}{3\sqrt{3}} \right)$$

$$= \frac{\pi}{4} \left(\frac{24a^3 - 8a^3}{3\sqrt{3}} \right)$$

$$= \frac{\pi}{4} \times \frac{16a^3}{3\sqrt{3}}$$

$$= \frac{4\pi a^3}{3\sqrt{3}} u^3 \text{ as reqd.}$$

(11) a) Prove true for $n = 1$,
 $9^1 - 5^1 = 4$, which is divis. by 4.

Assume true for $n = k$, i.e. assume that $9^k - 5^k = 4P$
for some integer P .

Prove true for $n = k+1$, i.e. prove that $9^{k+1} - 5^{k+1} = 4Q$
for some integer Q .

Now, $9^{k+1} - 5^{k+1} = 9 \times 9^k - 5 \times 5^k$





$$= 9 \times (4P + 5^k) - 5 \times 5^k \text{ from above}$$

$$= 36P + 4 \times 5^k$$

$$= 4(9P + 5^k)$$

$$= 4Q \text{ since } 9P + 5^k \text{ is integral.}$$

Since the result is true for $n = 1$, then it must be true for $n = 1 + 1 = 2$, then $n = 2 + 1 = 3$ and so on for all positive integers n .



c) $V = \pi \int_0^{\frac{\pi}{3}} \tan x dx$
 $= \pi \times \frac{\pi}{12} \left(\tan 0 + 4 \tan \frac{\pi}{12} + 2 \tan \frac{2\pi}{12} + 4 \tan \frac{3\pi}{12} + \tan \frac{\pi}{3} \right)$
 $= \frac{\pi^2}{12} \times 7.9585$
 $= 6.5 u^3$ (1 dec.)